

# Exome Sequencing and Genome-Wide Linkage Analysis in 17 Families Illustrate the Complex Contribution of *TTN* Truncating Variants to Dilated Cardiomyopathy

Nadine Norton, PhD; Duanxiang Li, MD, MS; Evadnie Rampersaud, PhD; Ana Morales, MS, CGC; Eden R. Martin, PhD; Stephan Zuchner, MD; Shengru Guo, MS; Michael Gonzalez, BSc; Dale J. Hedges, PhD; Peggy D. Robertson, PhD; Niklas Krumm, BA; Deborah A. Nickerson, PhD; Ray E. Hershberger, MD; on behalf of the National Heart, Lung, and Blood Institute GO Exome Sequencing Project and the Exome Sequencing Project Family Studies Project Team

**Background**—Familial dilated cardiomyopathy (DCM) is a genetically heterogeneous disease with >30 known genes. *TTN* truncating variants were recently implicated in a candidate gene study to cause 25% of familial and 18% of sporadic DCM cases.

**Methods and Results**—We used an unbiased genome-wide approach using both linkage analysis and variant filtering across the exome sequences of 48 individuals affected with DCM from 17 families to identify genetic cause. Linkage analysis ranked the *TTN* region as falling under the second highest genome-wide multipoint linkage peak, multipoint logarithm of odds, 1.59. We identified 6 *TTN* truncating variants carried by individuals affected with DCM in 7 of 17 DCM families (logarithm of odds, 2.99); 2 of these 7 families also had novel missense variants that segregated with disease. Two additional novel truncating *TTN* variants did not segregate with DCM. Nucleotide diversity at the *TTN* locus, including missense variants, was comparable with 5 other known DCM genes. The average number of missense variants in the exome sequences from the DCM cases or the ≈5400 cases from the Exome Sequencing Project was ≈23 per individual. The average number of *TTN* truncating variants in the Exome Sequencing Project was 0.014 per individual. We also identified a region (chr9q21.11-q22.31) with no known DCM genes with a maximum heterogeneity logarithm of odds score of 1.74.

**Conclusions**—These data suggest that *TTN* truncating variants contribute to DCM cause. However, the lack of segregation of all identified *TTN* truncating variants illustrates the challenge of determining variant pathogenicity even with full exome sequencing. (*Circ Cardiovasc Genet.* 2013;6:144-153.)

**Key Words:** dilated cardiomyopathy ■ exome ■ genetics ■ genome-wide analysis ■ human

Whole exome-sequencing technologies are rapidly enabling the identification of novel rare variants in patients with cardiomyopathy, but assigning pathogenicity remains challenging. Truncating variants in *TTN* were recently observed in 25% of familial dilated cardiomyopathy (DCM) cases.<sup>1</sup> DCM is genetically heterogeneous with rare variants in >30 disease genes, including *TTN*, previously indicated to cause DCM.<sup>2,3</sup> Before the recent publication of *TTN* contributing to a major fraction of genetic DCM, the fraction of cases attributable to any single gene ranged from <0.5% to ≈6% per disease gene.<sup>4</sup>

## Clinical Perspective on p 153

Discovery and incorporation into clinical tests of a single gene accounting for a large fraction of DCM cases could be helpful

for presymptomatic diagnosis in at-risk family members, but the clinical translation of this finding is confounded by several factors. First, despite a significant excess of truncating variants in DCM cases, these variants also occur in ≈3% of controls.<sup>1</sup> This is not unusual in complex trait analysis, where common genetic variants occur more frequently but not exclusively in cases compared with controls and increase disease risk or susceptibility. However, in the context of DCM, which has been categorized primarily as a rare-variant mendelian disease with marked locus and allelic heterogeneity,<sup>5</sup> it is essential to know which truncating variants are pathogenic. Second, with >300 exons and >34 000 amino acids, *TTN* has the largest coding sequence in the genome, and the majority of the general population will have at least 1 rare (defined as a mean allele frequency <0.5%) missense or truncating variant

Received June 28, 2012; accepted January 31, 2013.

From the Cardiovascular Division, Department of Medicine (N.N., D.L., A.M., R.E.H.), and Hussman Institute for Human Genomics (E.R., E.R.M., S.Z., S.G., M.G., D.J.H.), University of Miami Miller School of Medicine, Miami, FL; and Department of Genome Sciences, University of Washington, Seattle (P.D.R., N.K., D.A.N.).

The online-only Data Supplement is available at <http://circgenetics.ahajournals.org/lookup/suppl/doi:10.1161/CIRCGENETICS.111.000062/-/DC1>.

Correspondence to Ray E. Hershberger, MD, Dorothy M. Davis Heart and Lung Research Institute, Room 304 DHLRI, 473 W 12th Ave, Columbus, OH 43210. E-mail Ray.Hershberger@osumc.edu

© 2013 American Heart Association, Inc.

*Circ Cardiovasc Genet* is available at <http://circgenetics.ahajournals.org>

DOI: 10.1161/CIRCGENETICS.111.000062

at this locus. Next-generation sequencing now allows rapid variation analysis of the *TTN* gene despite its size. However, it relies on an economy of scale, and at a cost similar to that of sequencing *TTN* alone, next-generation sequencing can be used to sequence the entire coding sequence of the genome. This allows DCM patients to be screened for sequence variants in *TTN* in parallel with all other known DCM genes and the rest of the coding genome, leading to the third issue: The recent study of *TTN* truncating variants in DCM<sup>1</sup> used a custom next-generation sequencing panel specific for *TTN*, and therefore neither the role of genetic variants in other known DCM genes nor segregating variants in novel DCM genes could be assessed. Detailed examination of *TTN* truncating variants in the context of all coding variants in known and potentially novel DCM genes is needed to assess variant pathogenicity.

In this study, we used exome sequence data from 17 families, each with  $\geq 3$  members affected with DCM, in an effort to identify a genetic cause of DCM, because each family proband was negative for mutations in the coding regions of 16 DCM genes, as previously reported.<sup>6–11</sup> From these exome sequences, we identified several families with *TTN* truncating variants. To more carefully assess *TTN* variants as a cause of DCM, we constructed a linkage map of common informative single-nucleotide variants (SNVs) from our exome data in all 17 families and performed linkage analysis across the genome. We hypothesized that if *TTN* variants were causative of 25% of DCM, we would observe a significant combined logarithm of odds [LOD] score across our 17 families at this locus compared with the rest of the genome. Second, we evaluated all other unbiased rare variations in the exome data to identify putative DCM causative variants in each family. We describe the variants identified in *TTN* in the context of other top-ranking variants across the exome sequence of each family. Third, we examined the nucleotide diversity at the *TTN* locus in the 5400 exome sequences available from the Exome Variant Server<sup>12</sup> to determine whether the large amount of variation within this gene is accounted for by size or whether the *TTN* locus is more genetically diverse than other known DCM genes.

## Materials and Methods

### Subjects

Written informed consent was obtained from all subjects, and the Institutional Review boards at the Oregon Health and Science University and the University of Miami approved the study. The investigation included 17 families, each with  $\geq 3$  members affected with DCM and with each proband already known to be point mutation negative for 16 known DCM genes.<sup>6–11</sup> Genomic DNA was extracted from whole blood according to a standard salting-out procedure, as previously reported.<sup>6–11</sup>

### Linkage Analysis

Two-point and multipoint parametric linkage analyses were performed with the Merlin software<sup>13</sup> program. We assumed an affected-only model with a disease allele frequency of 0.0001 and penetrance of 0.9. In addition to traditional LOD scores, a heterogeneity LOD score resulting from a test of linkage in the presence of genetic heterogeneity was calculated. A genome-wide linkage map of common informative markers was constructed by identifying all SNVs present from the exome data that overlap with known SNVs in 60 unrelated Europeans from the International HapMap. SNVs with minor allele frequency <1% or mendelian errors within the HapMap were excluded. The remaining markers were then pruned with the PLINK software<sup>14</sup> using

pairwise  $r^2 < 0.1$  in sliding windows of 50 SNVs, moving in intervals of 5 SNVs. This resulted in a final exome-wide marker set of 4601 SNVs. Marker allele frequencies for linkage analysis were determined by the frequency of each SNV in the Exome Sequencing Project (ESP) in the relevant ethnically matched population, either European ( $n=3499$  individuals) or African American ancestry ( $n=1864$  individuals).

### Exome Sequencing and Analysis

Exome sequencing was performed at the University of Washington Genome Science Center across 17 families (48 individuals) with NimbleGen V2 in solution capture and Illumina HiSeq. Sequences were aligned with the Burrows-Wheeler Aligner,<sup>15</sup> and realignment and single-nucleotide and insertion-deletion variants were called with GATK version 1.4 at the Hussman Institute for Human Genomics. Vcf files were then imported into an in-house database, Genomes Management Application (GEMapp), to facilitate storage, variant annotation, querying, and analysis. In addition to our exome data, GEMapp was used to store transcriptome data from the left ventricle of 4 unrelated individuals, 2 with DCM and 2 unaffected individuals, as previously published.<sup>11</sup> This allowed us to filter variants mapped to genes expressed in heart tissue.

Using GEMapp, we queried each family to determine putative disease-causing variants that met the following criteria: variants with read depth  $\geq 5$  and quality scores  $\geq 40$ ; variants that were missense, nonsense, splice site, or a coding insertion or deletion; variants shared across all affected members of a family; variants with a frequency <0.5% in 5400 exomes from the Exome Variant Server (EVS); variants that have either a Phastcons<sup>16</sup> score >0.4 or a Genomic Evolutionary Rate Profiling<sup>17</sup> score >2; and variants with expression in our heart transcriptome data set with Reads Per Kilobase per Million mapped reads >3. These criteria were defined from our previous work on 197 variants in known DCM genes published as disease causing and that was analyzed by our group.<sup>18</sup> We also excluded filtered variants that were present in all 48 exomes and variants occurring in >1 family that did not segregate with disease status in at least 1 other family.

### Copy Number Variation

Copy number variation in 48 DCM exomes was also assessed by the Structural Variant Working Group at the University of Washington using CoNIFER (Copy Number Inference From Exome Reads; <http://conifer.sourceforge.net/>).<sup>19</sup> A total of 200 non-DCM exomes and 48 DCM exomes were used. Singular value decomposition transformation was used to remove systematic bias, removing 8 components. The final singular value decomposition -ZRPKM signal was then smoothed, and the duplication/deletion break points were found using a threshold of  $\pm 1.5$  singular value decomposition -ZRPKM.

### Sanger Sequencing Validation

All variants passing filter criteria and occurring within *TTN* were validated with Sanger sequencing and run on a 3130xl, as previously published.<sup>11</sup> Primer sequences are shown in Table I in the online-only Data Supplement. Additional DNA samples ( $n=29$ ) from affected and unaffected family members were also sequenced for these variants.

### Nucleotide Diversity

A total of 316 exons in *TTN* were targeted in our exome sequence. To assess the genetic variation at this locus, accounting for the large amount of coding sequence, we used the normalized number of variant sites,  $\theta$ , as a measure of nucleotide diversity across the 5379 exomes available from the Exome Variant Server.  $\theta$  was calculated as described by Cargill et al<sup>20</sup> for all coding sequence and separately for both missense and truncating variants.

## Results

### Exome-wide Linkage Analysis Across 17 Families With DCM

The maximum LOD score across the genome and the LOD score at the *TTN* locus are shown for each family in Table

**Table 1. Genome-wide Linkage and Rare Variant Exome Sequence Analysis Summary by Family**

| Family   | MLOD  | hg19 Region              | <i>TTN</i> MLOD | Filtered Exome Variants ( <i>TTN</i> Variants), n |
|--|-------|--------------------------|-----------------|---|
| Pedigrees with segregating <i>TTN</i> truncating variants          |       |                          |                 |   |
| A  | 1.178 | chr19:57802806-58058739  | 0.722           | 1 ( <i>TTN</i> Arg13527stop)                      |
| B  | 0.899 | chr11:1782594-5625847    | 0.725           | 6 ( <i>TTN</i> Ser19378stop, Ile2686Val)          |
| C  | 0.902 | chr6:35477032-42666164   | 0.034           | 12 ( <i>TTN</i> IVS 275+2 T>A)                    |
| D  | 0.602 | chr18:6890434-7017322    | 0.444           | 34 ( <i>TTN</i> IVS 275+2 T>A, Gly29127Arg)       |
| E  | 0.301 | N/A*                     | 0.283           | 28 ( <i>TTN</i> Val28259SerfsX22)                 |
| F  | 0.301 | N/A*                     | 0.236           | 24 ( <i>TTN</i> Lys28880AsnfsX8)                  |
| G  | 0.601 | chr11:290816-6891605     | 0.544           | 24 ( <i>TTN</i> Arg31175stop)                     |
| Other pedigrees without segregating <i>TTN</i> truncating variants |       |                          |                 |   |
| Family 8   | 0.901 | chr10:62863518-64597506  | -2.173          | 10  |
| Family 9   | 0.474 | chr5:137754695-137426447 | N/A             | 16  |
| Family 10  | 0.601 | chr11:62863518-64597506  | N/A             | 18  |
| Family 11  | 0.301 | N/A*                     | 0.296           | 22  |
| Family 12  | 0.602 | chr22:29456733-35660875  | N/A             | 29  |
| Family 13  | 0.301 | N/A*                     | -0.347          | 36  |
| Family 14  | 0.301 | N/A*                     | -0.305          | 43  |
| Family 15  | 1.242 | chr10:99504630-100219374 | N/A             | 43  |
| Family 16  | 0.301 | N/A*                     | -1.238          | 51  |
| Family 17  | 0.301 | N/A*                     | -0.66           | 80  |

MLOD indicates multipoint logarithm of odds (LOD) score across genome; N/A, not applicable because single-nucleotide variants at *TTN* locus were uninformative for linkage; and *TTN* MLOD, multipoint LOD score at *TTN* locus, number filtered exome variants refers to the number of top-ranking variants from exome pipeline with annotation of all top-ranking truncating and missense variants in *TTN*. Pedigrees A–G and 8–14 had 3 and pedigrees 15–17 had 2 affected family members, who underwent exome sequencing.

\*Not applicable in that >1 region had the same MLOD score.

1. In those families with *TTN* truncating variants identified by exome sequencing analysis, the maximum LOD score achieved across the genome for each family was comparable to the LOD score at the *TTN* locus (Table 1). The highest multipoint peak within the genome fell in the region spanning chromosome 9q21.11–q22.31 (hg19:71,862,987–95,840,256), producing a heterogeneity LOD score of 1.74. Overall, in the 17 families, the *TTN* locus was the second highest (heterogeneity LOD=1.59; Table 2).

### Exome Sequence Analysis From 17 Families With DCM

Our criteria for putative DCM variants identified in the exome sequences were based on defined criteria (Methods) and as previously described.<sup>11</sup> The number of shared variants meeting these criteria present in each family ranged from 1 to 80 (average, 28.1; median, 24). We had previously reported that of the 197 variants already published as causative of DCM, 16% were present in 2400 exomes from the ESP, and of those with functional data (and therefore presumed to be pathogenic variants of very low frequency), the median frequency in the ESP population was 0.04%.<sup>18</sup> When this maximum 0.04% frequency criterion was applied to the present exome analysis, the number of shared filtered variants per family ranged from 1 to 49 (average, 16.8; median, 15). Copy number variant analysis using exome data did not identify any shared rare variants (frequency <1% in the ESP data set) across these families.

A total of 6 *TTN* truncating variants (2 frameshift, 3 nonsense, and 1 splice variant that occurred in 2 DCM families) were identified among the filtered candidates in 7 of our 17 families (41%; Table 3). Our approach to identifying which of these truncating variants were likely disease causing within these families was to genotype them in additional DNA samples in the extended families when possible to assess segregation of the variant with disease; we also consider them in the context of the additional shared variants in our filtered lists for each family. We further observed from our linkage data that those families with highly negative LOD scores at the *TTN* locus had no *TTN* truncating variants that passed our exome analysis filtering criteria. In addition, we screened against presence in the 1000 Genome data (which are independent of the EVS data set). None of the 6 truncating variants were present in this data set.

**Table 2. Top 5 Positive Linkage Regions in 17 Families With Dilated Cardiomyopathy**

| chr | hg19 Start  | hg19 End    | Maximum HLOD |
|-----|-------------|-------------|--------------|
| 9   | 71 862 987  | 95 840 256  | 1.743        |
| 2   | 174 128 513 | 215 820 013 | 1.588        |
| 6   | 311 938     | 6 318 795   | 1.304        |
| 3   | 336 508     | 16 268 974  | 1.11         |
| 5   | 94 994 339  | 138 456 815 | 1.072        |

HLOD indicates heterogeneity logarithm of odds score. *TTN* locus is the second row of the table.

**Table 3. TTN Filtered Variants Identified in Exome Analysis and Validated With Sanger Sequencing That Segregated in Families With Dilated Cardiomyopathy**

| Hg19 Position  | Family Identification | Variant Function Class | PhastCons | GERP | Grantham | Additional Filtered Variants in Family | MAF % (EA) | MAF % (AA) | cDNA           | Protein          |
|----------------|-----------------------|------------------------|-----------|------|----------|--|------------|------------|----------------|------------------|
| Truncating     |                       |                        |           |      |          |  |            |            |                |                  |
| chr2:179481235 | A: F128               | Stop-gained            | 1.00      | 3.90 | N/A      | 0 (0)                                  | 0          | 0.03       | c.40579C>T     | Arg13527stop     |
| chr2:179447693 | B: F533               | Stop-gained            | 1.00      | 5.02 | N/A      | 4 (2)                                  | 0          | 0          | c.58133C>G     | Ser19378stop     |
| chr2:179424036 | C: F27                | Splice-5               | 0.99      | 5.61 | N/A      | 11 (8)                                 | 0          | 0          | N/A            | IVS 275+2 T>A    |
| chr2:179424036 | D: F40                | Splice-5               | 0.99      | 5.61 | N/A      | 32 (17)                                | 0          | 0          | N/A            | IVS 275+2 T>A    |
| chr2:179413874 | E: F2B                | Frameshift             | N/A       | N/A  | N/A      | 27 (14)                                | 0          | 0          | c.84774insT    | Val28259SerfsX22 |
| chr2:179411904 | F: F19                | Frameshift             | N/A       | N/A  | N/A      | 23 (18)                                | 0          | 0          | c.86640delAGAA | Lys28880AsnfsX8  |
| chr2:179400115 | G: F35                | Stop-gained            | 1.00      | 4.64 | N/A      | 23 (10)                                | 0          | 0          | c.93523C>T     | Arg31175stop     |
| Missense       |                       |                        |           |      |          |  |            |            |                |                  |
| chr2:179635998 | B: F533               | Missense               | 0.98      | 2.44 | 29       | 4 (2)                                  | 0          | 0          | c.8056A>G      | Ile2686Val       |
| chr2:179410975 | D: F40                | Missense               | 1.00      | 5.66 | 125      | 32 (17)                                | 0          | 0.06       | c.87379G>A     | Gly29127Arg      |

MAF% indicates percent minor allele frequency from 5379 exomes in the Exome Variant Server (EVS), given in European American (EA) and African American (AA) populations. The number of additional filtered variants in other genes identified in each family with a TTN variant are given; the number of variants under more stringent filtering criteria of <0.05% in the EVS is given in parentheses. Conservation scores are given as PhastCons and GERP.

**Clinical Characteristics Families With DCM and Segregating TTN Variants**

The pedigree structures are shown (Figure) and the clinical characteristics of relevant family members are provided (Table 4) of the DCM families with segregating TTN variants (Table 1).


**Family A**

Family A had DNA samples available for 3 additional members. Sanger sequencing showed that all 6 family members were heterozygous for the truncating variant. Subject III.3, a woman who carried the TTN variant, died at 69 years of age with mild systolic dysfunction (ejection fraction, 42%) but without left ventricular enlargement, having suffered a myocardial infarction in her 50s, and thus confounding assessment of whether

the TTN variant, the myocardial infarction, or both contributed to her systolic dysfunction. Two subjects (IV.3, V.1), both mutation carriers in their 20s, had no evidence of DCM.

**Family B**

In addition to the 3 samples that had exome sequencing, family B had DNA samples available from 6 other family members. Sanger sequencing confirmed the nonsense variant as present in all affected family members. A female obligate carrier (II.2) died of cancer at 76 years of age without a cardiovascular history. Another female obligate carrier (II.5) had no cardiovascular history at 70 years of age. A male who carried the TTN variant at 69 years of age (II.6) had only borderline systolic dysfunction without left ventricular enlargement.



**Figure.** Pedigrees of families with dilated cardiomyopathy (DCM). Squares represent male subjects; circles represent female subjects. Diagonal lines mark deceased individuals. Solid symbols denote DCM. Gray symbols represent any cardiovascular abnormality. Open symbols represent unaffected individuals or individuals with no data available for analysis. The presence or absence of the family’s TTN truncating variant is indicated by a + or – symbol, respectively; obligate carriers are noted in parenthesis as (+) and unknown zygosity as (?). Individuals who underwent exome sequencing are denoted as (exome).

Downloaded from http://circgenetics.ahajournals.org/ by guest on June 24, 2017

Table 4. Clinical Characteristics

| Subject                   | Age at Diagnosis, Screening or Family History Information, y | Sex | DCM | ECG/Arrhythmia    | LVEDD, mm (Z score) | LV Septum,                   |               | EF or FS, % | TTN Truncating Mutation Present | Other Mutation? (Genotype)         | Comment   |
|---------------------------|--|-----|-----|-------------------|---------------------|------------------------------|---------------|-------------|---------------------------------|------------------------------------|---|
|                           |  |     |     |                   |                     | Posterior Wall Thickness, mm | Thickness, mm |             |                                 |                                    |   |
| Pedigree A: Arg13527X     |  |     |     |                   |                     |                              |               |             |                                 |                                    |   |
| IV-2                      | 42   | F   | Yes | PVCs              | 56, NA              | NA                           | NA            | 20          | Yes                             |                                    |   |
| III-1                     | 43   | F   | Yes |                   | 54 (2.44)           | NA, 11                       | NA, 11        | 40          | Yes                             |                                    |   |
| III-3                     | 69   | F   | No  | Anterior MI       | 54, NA              | 11, 10                       | 11, 10        | 42          | Yes                             |                                    | CAD diagnosis at age 69   |
| III-5                     | 51   | M   | Yes | PVCs, AF          | 67 (4.04)           | NA                           | NA            | 35          | Yes                             |                                    | history: paroxysmal atrial tachycardia and syncope  |
| IV-3                      | 27   | F   | No  |                   |                     |                              |               |             | Yes                             |                                    | records documenting normal echocardiogram   |
| V-1                       | 23   | F   | No  |                   |                     |                              |               |             | Yes                             |                                    |   |
| Pedigree B: Ser19378X     |  |     |     |                   |                     |                              |               |             |                                 |                                    |   |
| III-3                     | 29   | M   | Yes | PVCs              | 56 (1.54)           | 10, 10                       | 10, 10        | 47          | Yes                             | TTN Ile2640Val                     | history: died at 76 from cancer   |
| II-2                      | NA   | F   | No  |                   |                     |                              |               |             | Obligate carrier                | obligate carrier of TTN Ile2640Val |   |
| II-3                      | 66   | M   | No  |                   |                     |                              |               |             | No                              |                                    | history: normal cardiac screening   |
| II-5                      | 70   | F   |     |                   |                     |                              |               |             | Obligate carrier                | obligate carrier of TTN Ile2640Val | history: heart attack at age 50–60.   |
| II-6                      | 64   | M   | No  |                   | 56 (1.42)           | 9, 8                         | 9, 8          | 49.5        | Yes                             | TTN Ile2640Val                     | Borderline systolic dysfunction   |
| II-7                      | 62   | F   | No  |                   | 42 (–1.15)          | 10, 10                       | 10, 10        | 39 (FS)     | No                              |                                    |   |
| III-1                     | 43   | M   | No  | ICD               |                     |                              |               |             | Yes                             | TTN Ile2640Val                     | CAD diagnosis at age 43   |
| III-2                     | 47   | F   | Yes | PVCs, bigeminy    | 64 (5.21)           | 8, 6.8                       | 8, 6.8        | 21          | Yes                             | TTN Ile2640Val                     |   |
| III-4                     | 36   | M   | Yes | ICD, PVCs         | 64 (3.43)           | 9, 10                        | 9, 10         | 24.5        | Yes                             | TTN Ile2640Val                     |   |
| Pedigree C: IVS 275+2 T>A |  |     |     |                   |                     |                              |               |             |                                 |                                    |   |
| III-1                     | 37   | F   | Yes | NSSTT             | 68 (5.6)            | NA                           | NA            | 40          | Yes                             |                                    | Right deltoid muscle biopsy attributable to history of muscle weakness revealed congenital type I fiber predominance or chronic neurogenic atrophy with reinnervation.<br>History: died at 65 from cancer |
| I-2                       | NA   | F   |     |                   |                     |                              |               |             | Obligate carrier                |                                    |   |
| II-2                      | 52   | F   | Yes | NSSTT             | 57 (3.29)           | 11, 11                       | 11, 11        | 22          | Yes                             |                                    |   |
| II-3                      | 59   | M   | No  |                   |                     |                              |               |             | Obligate carrier                |                                    | Death certificate: VT, CAD  |
| III-2                     | 39   | M   | Yes | PVCs, LAE, NSSTT  | 73 (4.69)           | 10, 10                       | 10, 10        | 32          | Yes                             |                                    |   |
| Pedigree D: IVS 275+2 T>A |  |     |     |                   |                     |                              |               |             |                                 |                                    |   |
| II-4                      | 39   | M   | Yes | AF, cardioversion | 66 (3.58)           | 11, 11                       | 11, 11        | 39          | Yes                             | TTN Gly29127Arg                    |   |
| II-3                      | 49   | F   | Yes | PVCs, PACs, MI    | 58 (3.56)           | 10, 10                       | 10, 10        | 25 (FS)     | Yes                             | TTN Gly29127Arg                    |   |
| II-5                      | 46   | F   | Yes | NSSTT             | 53 (2.26)           | 8, 8                         | 8, 8          | 46          | Yes                             | TTN Gly29127Arg                    |   |
| III-1                     | 32   | M   | No  | RBBB, MI          | 51 (–0.07)          | 8, 8                         | 8, 8          | 41          | Yes                             |                                    |   |

(Continued)



Table 4. (Continued)

|                              |    |   |     |                         |             |       |         |     |                  |  |
|------------------------------|----|---|-----|-------------------------|-------------|-------|---------|-----|------------------|--|
| Pedigree E: Val28259SerfsX22 |    |   |     |                         |             |       |         |     |                  |  |
| II-3                         | 14 | M | yes |                         | 75 (6.30)   | 5, 8  | 13 (FS) | yes |                  | Heart transplantation  |
| I-1                          | 44 | M | yes | LAHB, MI, NSSTT         | 62 (3.06)   | 10, 9 | 25      | yes |                  |  |
| I-2                          | 42 | F | no  | Normal                  | 55 (2.32)   | 8, 9  | 60      | no  | MYBPC3 Ala833Thr |  |
| II-1                         | 20 | M | no  | Normal                  | 60 (2.28)   | 10, 9 | 60      | yes |                  |  |
| II-2                         | 22 | M | yes | AF, ICD                 | 63.5 (3.01) | 10, 7 | 20      | yes | MYBPC3 Ala833Thr | Heart transplantation  |
| Pedigree F: Lys28880AsnfsX8  |    |   |     |                         |             |       |         |     |                  |  |
| III-1                        | 16 | M | yes | AF, NSSTT               |             |       | 21      | yes |                  | Cardiomegaly (CXR); heart transplantation. Died post-transplant, age 21. |
| II-1                         | 47 | M | no  | Normal                  | 52 (0.28)   | 12, 9 | 83      | no  |                  |  |
| II-2                         | 45 | F | yes | NSSTT                   | 51 (1.78)   | NA    | 48      | yes |                  |  |
| II-3                         | 52 | M | yes | AF, NSCD, cardioversion | 50 (NA)     | 8, 9  | 35      | yes |                  | Death from noncardiovascular cause                                       |
| II-5                         | 52 | M | no  | 1AVB                    | 50 (-0.22)  | 9, 10 | 77      | no  |                  |  |
| II-6                         | 41 | F | no  | MI, NSSTT               | 55 (2.97)   | 6, 8  | 68      | no  |                  |  |
| III-2                        | 20 | M | yes | ICD, tachycardia        | 60 (2.55)   | 8, 6  | 38      | yes |                  | Heart transplantation  |
| III-3                        | 25 | F | no  | Normal                  | 51 (0.53)   | 7, 7  | 47      | yes |                  |  |
| Pedigree G: Arg31175X        |    |   |     |                         |             |       |         |     |                  |  |
| II-1                         | 52 | M | yes |                         | 56 (NA)     | NA    | 10      | yes |                  | By history: no known heart problems                                      |
| II-2                         |    | M |     |                         |             |       |         |     | obligate carrier |  |
| II-4                         | 49 | M | yes | LAE, tachycardia, NSSTT | 70 (5.04)   | 12, - | FS 17   | yes |                  |  |
| III-1                        | 35 | M | yes | LAE, tachycardia, NSSTT | 69 (5.21)   | 10, 9 | 20      | yes |                  |  |

1AVB indicates first-degree atrioventricular block; AF, atrial fibrillation; DCM, dilated cardiomyopathy; EF, ejection fraction; FS, fractional shortening; ICD, implantable cardiac defibrillator; LAE, left atrial enlargement; LAHB, left anterior hemiblock; LV, left ventricle; LVEDD, left ventricular end-diastolic dimension; MI, myocardial infarction pattern; NA, not available; NSCD, nonspecific conduction delay; NSR, normal sinus rhythm; NSSTT, nonspecific ST-T changes; PAC, premature atrial contraction; PVC, premature ventricular contraction; and RBBB, right bundle-branch block.

**Family C**

Only 3 DNA samples were available for this family and were used for exome sequencing. Subject III.3, who died of DCM, was an identical twin by family history and thus may have been an obligate carrier. Another obligate carrier (III.3) had no known DCM but according to the death certificate died of ventricular tachycardia and coronary artery disease. None of the additional 12 variants passing filtering criteria (8 under more stringent filtering of population frequency <0.05% in the ESP exome data set) occurred in known DCM or other cardiomyopathy-associated genes.

**Family D**

This family of European ancestry carried the same *TTN* splice variant identified in family C. Of the additional 32 variants also identified as putative disease causing in this family, only 1 variant occurred in a known DCM gene, a missense variant in *TTN*, chr2:179,410,975, NM\_133378.4, Gly29127Arg.

**Family E**

This family carried a single base insertion in *TTN*, resulting in a frameshift mutation. Of the additional 26 segregating variants also identified in this family, the only variant in a gene with a reported association with hypertrophic cardiomyopathy occurred in *SOS1*.<sup>21</sup> One of the affected children (II.2) carried a rare variant in *MYBPC3* inherited from his mother (I.2), previously reported by us as likely disease causing<sup>9</sup> but suggested to be of unknown significance on the basis of a subsequent study<sup>22</sup>; we also note that this variant is present at a frequency of 0.12% in the EVS, making it more common than many DCM rare variants.<sup>18</sup>

**Family F**

A 4-bp deletion in *TTN* resulted in a frameshift mutation. DNA was available from 1 additional affected member and 3 unaffected family members, and the *TTN* variant was confirmed to be present in all 4 affected members by Sanger sequencing and was not present in the 3 unaffected members. None of the other 23 segregating variants identified occurred in known cardiomyopathy-associated genes.

**Family G**

No DNA samples beyond those used for exome sequencing were available to assess segregation in this family. Of the additional 23 variants that segregated, none occurred in other cardiomyopathy-associated genes.

**Segregating Missense Variants at the *TTN* Locus**

We also considered the implication of segregating *TTN* missense variants passing our exome-filtering pipeline as a class of variants that were not discussed in the recent *TTN* study.<sup>1</sup> Determination of pathogenicity of these variants will be extremely challenging because of the large number of coding exons. In the 5400 ESP exome data sets, the average number of *TTN* missense variants per individual was 23.3, ranging 6 to 55. These results were comparable to those observed in the exome sequences of our DCM families, with the average number of *TTN* missense variants per DCM individual at 22.75 (range, 11–43). Application of our DCM filtering criteria to missense variants in the 5400

ESP exomes (frequency <0.5% and either a PhastCons score >0.4 or a GERP score of >2) resulted in an average of 1.91 missense variants per individual (range, 0–23). Five *TTN* missense variants passed our exome-filtering approach (that segregated with all individuals affected with DCM in a family): 1 each in 2 DCM families who also had segregating truncating variants (Table 3) and the others in 2 families, each with high-quality candidates in known cardiomyopathy genes, so they were not further prioritized. The average number of *TTN* missense variants without regard to sharing, that is, an analysis of only 1 individual from each of the 17 families, a less stringent approach and similar to the analysis of all missense *TTN* variants conducted for the EVS, was 1.88.

**Nonsegregating, Truncating Variants**

Given the observed excess of *TTN* truncating variants in both familial and in sporadic DCM versus controls,<sup>1</sup> we thought it also relevant to report the number of nonsegregating truncating variants in *TTN* identified in the exome sequences of these 48 individuals with DCM, because they may be potential susceptibility variants. We observed 2 nonsegregating *TTN* truncating variants that were validated with Sanger sequencing. First, a C insertion at hg19 chr2:179,426,992, generating a frameshift in 1 of 3 family members with DCM who underwent exome sequencing (family 14; Table 1) and a nonsense variant at hg19 chr2:179,605,218, NM\_003319.4 Gln3885stop in 2 of 3 family members (family 17; Table 1). Neither variant was observed in the 5400 ESP exome sequences or in the 1000 Genomes data, making them potential susceptibility variants. In the case of the frameshift variant, this family had already been shown to segregate a variant published as disease causing accompanied by functional data,<sup>6</sup> and in family 14 with the nonsense variant, a total of 43 segregating variants were identified by our exome-filtering pipeline (Table 1), none of which were in previously published cardiomyopathy genes.

**Nucleotide Diversity of *TTN***

The NimbleGen V2 in solution capture target included 315 discrete exons from 6 *TTN* transcripts (NM\_001256850.1, NM\_133432.3, NM\_133378.4, NM\_003319.4, NM\_133437.3, and NM\_133379.3), totaling 110 459-bp coding sequence. Our exome pipeline identified *TTN* truncating variants in 7 DCM families. This could simply be a result of the large number of exons. Hence, we investigated the nucleotide diversity at the *TTN* locus in a non-DCM population. The EVS contains annotated exome sequence from 5379 individuals at the *TTN* locus, totaling 2425 and 25 discrete missense and nonsense variants, respectively. Across 5379 EVS individuals, there are a total of 125 575 missense alleles and 77 nonsense alleles, averaging 23 and 0.014 per individual, respectively. We calculated the normalized number of variant sites,  $\theta$ ,<sup>20</sup> accounting for sample size and the number of coding bases in the EVS individuals at the *TTN* locus to be  $2.23 \times 10^{-3}$  and  $2.3 \times 10^{-5}$  for missense and nonsense variants, respectively. The same calculation across 5 other known DCM genes (*MYBPC3*, *TNNC1*, *TNNI3*, *MYH6*, and *TPMI*) in the EVS data gave results comparable to those observed in *TTN* (for missense variants,  $\theta$

ranged from  $6.9 \times 10^{-4}$  at *TPM1* to  $1.05 \times 10^{-3}$  at *TNNC1* and for truncating variants, from 0 at *TNNC1* and *TNNI3* to  $1.29 \times 10^{-4}$  for *MYBPC3*), suggesting that the excess of shared truncating variants in our DCM families is not attributable to the large number of exons alone and that nucleotide diversity at *TTN* is comparable to that of other known DCM genes.

## Discussion

This is the first independent replication study of *TTN* truncating variants as frequently involved in the pathogenesis of familial DCM. Herman et al<sup>1</sup> recently identified *TTN* truncating mutations in 25% of familial DCM and 18% of sporadic DCM, a significant excess compared with 3% of controls. The authors concluded that truncating mutations in *TTN* are a frequent cause of DCM because all previous reports of unselected patients with DCM of unknown cause ranged from <<1% to 5% to 8%.<sup>4</sup> However, 3% of controls in the Herman et al<sup>1</sup> study also were observed to have *TTN* truncating variants, suggesting that the interpretation of specific *TTN* truncating variant pathogenicity would be challenging, especially in simple cases. Analysis across 19 DCM families, segregating rare *TTN* truncating variants in the Herman et al<sup>1</sup> study, yielded a combined LOD score of 11.1, providing strong evidence that the truncating variants in those families were pathogenic. However, in that study, *TTN* was sequenced in isolation, so the relevance of the linkage evidence in the *TTN* region could not be compared with the rest of the genome.

We hypothesized that if rare variants in *TTN* indeed account for one quarter of familial DCM, this locus should also be detected using an unbiased genome-wide linkage approach across our 17 DCM families because they should be enriched for causative variants at this locus, especially because the DCM families in this study were selected for exome sequencing, because they were already known to be point mutation negative for 16 other known DCM genes<sup>6–11</sup> (except for 1 family segregating a previously described variant<sup>6</sup> in a gene attributing  $\approx 0.5\%$  of DCM). Genome-wide linkage analysis yielded the second most significant evidence of linkage at the *TTN* locus compared with other regions in the genome, which we interpret as evidence of the *TTN* locus in DCM pathogenesis.

Next, we identified those nonsense, missense, splice, and frameshift variants in the exome sequences meeting our filtering approach that included conservation and myocardial expression, which segregated with DCM affection status in each family. Seven of 17 families (41%) had segregating *TTN* truncating variants identified in their filtered exome variants. Using common informative SNVs within the exome sequences, we obtained a combined LOD score at the *TTN* locus for these 7 families of 2.99, and in each family, the maximum LOD score at the *TTN* locus was either the maximum LOD score achieved in that family across the whole exome or comparable to the maximum observed LOD score at any other locus. We interpret these data as replication of *TTN* truncating variants as frequently linked with DCM.

Despite the previous evidence<sup>1</sup> and our findings presented here, all of which collectively support the concept that *TTN* truncating variants are highly relevant for DCM pathogenesis, determining the pathogenicity of any specific variant remains extremely challenging. We interpret the 2 truncating variants

not shared by all affected family members in 2 families (families 14 and 17) as unlikely to be causative of DCM. We also note that in the 7 families in whom all those affected with DCM carried a truncating variant, some unaffected members at older ages also carried the same truncating variant. This observation confounds pedigree analysis, although it is consistent with reduced penetrance, which is commonly observed with familial DCM. Furthermore, the plethora of *TTN* missense variants observed in all individuals, whether from control or DCM cohorts, further complicates *TTN* variant interpretation. The available evidence from the ESP data has shown that most individuals will carry numerous *TTN* missense variants, some even very rare, and even if such variants segregate with DCM in a family, it may occur as a play of chance. This concept may also apply to truncating variants.

These issues raise 2 central questions of *TTN* biology in DCM. Which specific variants, whether truncating or missense, play a role in DCM pathogenesis? Do *TTN* variants include causative and risk alleles? Titin splicing and titin biology are exceedingly complex,<sup>23,24</sup> and penetrance is well known to be an incomplete and expressivity variable in familial DCM,<sup>4,5</sup> so it is possible if not likely that some *TTN* variants, whether missense or truncating, may also modulate penetrance and expressivity in DCM. Addressing these questions will require much larger DCM cohorts with detailed phenotypic data, ideally with knowledge of extended family structure (including presymptomatic DCM), genome-wide sequence data, and comprehensive, insightful analysis of the pathophysiological effects of *TTN* variants.

Although linkage analysis has been used less frequently in the genome-wide association study era, our study highlights the importance of coupling linkage information with sequence data. This provides us both a measure of evidence for a cumulative effect of rare variants, because linkage is not compromised by allelic heterogeneity (ie, multiple rare disease variants within a gene), and an assessment of the evidence in the context of the rest of the genome. Together, linkage analysis and sequencing provide complementary evidence that can improve the efficiency of gene discovery in sequencing studies.

We observed that 41% (7/17) of families carried *TTN* variants that segregated with DCM, which is higher than the 25% frequency observed in the Herman et al<sup>1</sup> study. This most likely resulted from a sample bias in our study because our families were already known to be point mutation negative for 16 other known DCM genes and thus were likely enriched for *TTN* variants.

We also examined *TTN* missense variants in our 17 families using an unbiased approach to exome analysis and additional data from >300 exomes with neurological phenotypes collated in our in-house database, GEMapp. Our calculation of nucleotide diversity in the ESP data set for this gene suggested that diversity at the *TTN* locus was comparable to other known DCM genes relative to the number of coding nucleotides. However, the >300 *TTN* exons resulted in a very large number of missense variants identified in both the exome sequence from individuals with DCM and individuals in the ESP data set. Two *TTN* missense variants from 2 DCM families, both also carrying *TTN* truncating variants, passed our exome-filtering criteria. We considered these 2 missense



variants (Gly29127Arg and Ile2685Val) of unknown significance because each met our stringent exome-filtering criteria and were not present in >300 other exomes in GEMapp but occurred in families B and D, who also had *TTN* truncating variants.

We also note that the most highly linked region in this study on chromosome 9q21.1-q22.31 did not contain any known DCM genes and that none of our filtered genes mapped to this region. The positive linkage at this region could be a chance finding. Alternatively, this could represent a region of the genome containing a novel DCM gene missed by our exome pipeline for 2 possible reasons. Our rare variant exome analysis approach is based on assumptions based on our previous work.<sup>18</sup> First, we assumed that causative rare variants were missense, nonsense, splice, or frameshift and that the allele frequency of these variants would be <0.5%. We note here that the majority of known (published) DCM variants are significantly less frequent than 0.5% in the general population,<sup>18</sup> but there are examples of known DCM variants with convincing functional data in which the variant frequency is very close to this cut point (*CSRP3* Trp4Arg variant<sup>25</sup> has a frequency of 0.35% in the European ancestry EVS data set). Although this variant would have been identified in our pipeline, it is possible that other pathogenic variants have frequencies slightly greater than this. Second, our genome-wide linkage approach could also have identified regions containing common susceptibility or modifying variants, again, that would not have been detected by our exome analysis approach. We note recent evidence of linkage to congenital heart defects and low atrial rhythm to this region,<sup>26</sup> suggesting the possibility of cardiovascular modifying variants located here.

In conclusion, our data show that *TTN* was the only gene with implicated rare variants that occurred in multiple DCM families and hence have replicated the previous finding<sup>1</sup> that *TTN* truncating variants contribute frequently to DCM pathogenesis. We reiterate that *TTN* analysis for DCM causation should be considered within the context of the genome. Whereas interpreting individual *TTN* truncating or missense variants will remain challenging because of the complexity of *TTN* biology, the availability of sequencing data from known DCM genes and variants at other exomic loci will assist in categorizing the pathogenicity of these variants.

### Acknowledgments

We thank all the family members who participated, without whom this study would not be a success. We would like to thank the National Heart, Lung, and Blood Institute GO ESP, which produced exome variant calls for comparison; the Lung GO Sequencing Project (HL-102923); the Women's Health Initiative (WHI) Sequencing Project (HL-102924); the Broad GO Sequencing Project (HL-102925); the Seattle GO Sequencing Project (HL-102926); and the Heart GO Sequencing Project (HL-103010). We would like to thank the Exome Sequencing Project Family Studies Project Team: Sek Kathiresan, Jay Shendure, Mike Bamshad, Weiniu Gan, Rebecca Jackson, Ani Manichaikul, Christopher Newton-Cheh, Debbie Nickerson, Stephen Rich, Jerry Rotter, and James Wilson.

### Sources of Funding

This work was supported by National Institutes of Health awards HL58626 (Dr Hershberger) and HL094976 (Dr Nickerson, Seattle Seq and the Stanley J. Glaser Award (Dr Norton).

### Disclosures

None.

### References

- Herman DS, Lam L, Taylor MR, Wang L, Teekakirikul P, Christodoulou D, et al. Truncations of titin causing dilated cardiomyopathy. *N Engl J Med*. 2012;366:619–628.
- Gerull B, Gramlich M, Atherton J, McNabb M, Trombitás K, Sasse-Klaassen S, et al. Mutations of *TTN*, encoding the giant muscle filament titin, cause familial dilated cardiomyopathy. *Nat Genet*. 2002;30:201–204.
- Gerull B, Atherton J, Geupel A, Sasse-Klaassen S, Heuser A, Frenneaux M, et al. Identification of a novel frameshift mutation in the giant muscle filament titin in a large Australian family with dilated cardiomyopathy. *J Mol Med*. 2006;84:478–483.
- Hershberger RE, Siegfried JD. Update 2011: clinical and genetic issues in familial dilated cardiomyopathy. *J Am Coll Cardiol*. 2011;57:1641–1649.
- Hershberger RE, Morales A, Siegfried JD. Clinical and genetic issues in dilated cardiomyopathy: a review for genetics professionals. *Genet Med*. 2010;12:655–667.
- Li D, Parks SB, Kushner JD, Nauman D, Burgess D, Ludwigsen S, et al. Mutations of presenilin genes in dilated cardiomyopathy and heart failure. *Am J Hum Genet*. 2006;79:1030–1039.
- Parks SB, Kushner JD, Nauman D, Burgess D, Ludwigsen S, Peterson A, et al. Lamin A/C mutation analysis in a cohort of 324 unrelated patients with idiopathic or familial dilated cardiomyopathy. *Am Heart J*. 2008;156:161–169.
- Hershberger RE, Parks SB, Kushner JD, Li D, Ludwigsen S, Jakobs P, et al. Coding sequence mutations identified in MYH7, TNNT2, SCN5A, CSRP3, LBD3, and TCAP from 313 patients with familial or idiopathic dilated cardiomyopathy. *Clin Transl Sci*. 2008;1:21–26.
- Hershberger RE, Norton N, Morales A, Li D, Siegfried JD, Gonzalez-Quintana J. Coding sequence rare variants identified in MYBPC3, MYH6, TPM1, TNNC1, and TNNI3 from 312 patients with familial or idiopathic dilated cardiomyopathy. *Circ Cardiovasc Genet*. 2010;3:155–161.
- Li D, Morales A, Gonzalez-Quintana J, Norton N, Siegfried JD, Hofmeyer M, et al. Identification of novel mutations in *RBM20* in patients with dilated cardiomyopathy. *Clin Transl Sci*. 2010;3:90–97.
- Norton N, Li D, Rieder MJ, Siegfried JD, Rampersaud E, Züchner S, et al. Genome-wide studies of copy number variation and exome sequencing identify rare variants in *BAG3* as a cause of dilated cardiomyopathy. *Am J Hum Genet*. 2011;88:273–282.
- Exome Variant Server, NHLBI Exome Sequencing Project (ESP). <http://evs.gs.washington.edu/EVS/>. Accessed August 8, 2012.
- Abecasis GR, Cherny SS, Cookson WO, Cardon LR. Merlin—rapid analysis of dense genetic maps using sparse gene flow trees. *Nat Genet*. 2002;30:97–101.
- Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet*. 2007;81:559–575.
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al; 1000 Genome Project Data Processing Subgroup. The Sequence Alignment/Map format and SAMtools. *Bioinformatics*. 2009;25:2078–2079.
- Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, Rosenbloom K, et al. Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. *Genome Res*. 2005;15:1034–1050.
- Cooper GM, Stone EA, Asimenos G, Green ED, Batzoglu S, Sidow A; NISC Comparative Sequencing Program. Distribution and intensity of constraint in mammalian genomic sequence. *Genome Res*. 2005;15:901–913.
- Norton N, Robertson PD, Rieder MJ, Züchner S, Rampersaud E, Martin E, et al; National Heart, Lung and Blood Institute GO Exome Sequencing Project. Evaluating pathogenicity of rare variants from dilated cardiomyopathy in the exome era. *Circ Cardiovasc Genet*. 2012;5:167–174.
- Krumm N, Sudmant PH, Ko A, O'Roak BJ, Malig M, Coe BP, et al; NHLBI Exome Sequencing Project. Copy number variation detection and genotyping from exome sequence data. *Genome Res*. 2012;22:1525–1532.
- Cargill M, Altshuler D, Ireland J, Sklar P, Ardlie K, Patil N, et al. Characterization of single-nucleotide polymorphisms in coding regions of human genes. *Nat Genet*. 1999;22:231–238.
- Roberts AE, Araki T, Swanson KD, Montgomery KT, Schiripo TA, Joshi VA, et al. Germline gain-of-function mutations in *SOS1* cause Noonan syndrome. *Nat Genet*. 2007;39:70–74.

22. Lakdawala NK, Funke BH, Baxter S, Cirino AL, Roberts AE, Judge DP, et al. Genetic testing for dilated cardiomyopathy in clinical practice. *J Card Fail.* 2012;18:296–303.
23. Guo W, Bharmal SJ, Esbona K, Greaser ML. Titin diversity: alternative splicing gone wild. *J Biomed Biotechnol.* 2010;753675:1–8.
24. Patel JR, Pleitner JM, Moss RL, Greaser ML. Magnitude of length-dependent changes in contractile properties varies with titin isoform in rat ventricles. *Am J Physiol Heart Circ Physiol.* 2012;302:H697–H708.
25. Knöll R, Hoshijima M, Hoffman HM, Person V, Lorenzen-Schmidt I, Bang ML, et al. The cardiac mechanical stretch sensor machinery involves a Z disc complex that is defective in a subset of human dilated cardiomyopathy. *Cell.* 2002;111:943–955.
26. van de Meerakker JB, van Engelen K, Mathijssen IB, Lekanne dit Deprez RH, Lam J, Wilde AA, et al. A novel autosomal dominant condition consisting of congenital heart defects and low atrial rhythm maps to chromosome 9q. *Eur J Hum Genet.* 2011;19:820–826.

### CLINICAL PERSPECTIVE

Rare variants that cause familial dilated cardiomyopathy (DCM) have been discovered in >30 genes but account for only 40% to 50% of cases. Exome sequencing, in which ≈19 000 coding genes in the human genome are sequenced simultaneously, provides a dramatically more powerful approach for DCM gene discovery. To this end, exome sequencing was undertaken in 3 or more affected members in 17 DCM families. Truncating rare variants in *TTN*, encoding the giant protein titin, were identified in 7 families and segregated with DCM and were not present in 5400 DNA exome sequences from the Exome Sequencing Project or the 1000 Genomes Project. A recent report suggested that rare *TTN* truncating variants cause up to 24% of familial dilated cardiomyopathy, but in that study, 3% of controls had *TTN* truncating variants, complicating assignment of pathogenicity to any specific *TTN* variant, and only *TTN* was sequenced (so in some cases mutations in other DCM genes that were not sequenced may have been relevant). The data from the present study strengthen the previous conclusion that *TTN* truncating variants are in general relevant for the pathogenesis of DCM: The genome-wide approach here excluded mutations in other known and novel DCM genes, and linkage analysis supported the pathogenicity of the *TTN* truncating variants in the 7 families. Nevertheless, 2 other *TTN* truncating variants did not segregate in 2 DCM families. Overall, our study suggests that *TTN* truncating variants contribute to the pathogenesis of DCM, but the lack of segregation of all identified *TTN* truncating variants illustrates the challenge of determining *TTN* variant pathogenicity even with exome sequencing.

**Exome Sequencing and Genome-Wide Linkage Analysis in 17 Families Illustrate the Complex Contribution of TTN Truncating Variants to Dilated Cardiomyopathy**

Nadine Norton, Duanxiang Li, Evadnie Rampersaud, Ana Morales, Eden R. Martin, Stephan Zuchner, Shengru Guo, Michael Gonzalez, Dale J. Hedges, Peggy D. Robertson, Niklas Krumm, Deborah A. Nickerson and Ray E. Hershberger  
on behalf of the National Heart, Lung, and Blood Institute GO Exome Sequencing Project and the Exome Sequencing Project Family Studies Project Team

*Circ Cardiovasc Genet.* 2013;6:144-153; originally published online February 15, 2013;  
doi: 10.1161/CIRCGENETICS.111.000062

*Circulation: Cardiovascular Genetics* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2013 American Heart Association, Inc. All rights reserved.  
Print ISSN: 1942-325X. Online ISSN: 1942-3268

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://circgenetics.ahajournals.org/content/6/2/144>

Data Supplement (unedited) at:

<http://circgenetics.ahajournals.org/content/suppl/2013/02/15/CIRCGENETICS.111.000062.DC1>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation: Cardiovascular Genetics* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Circulation: Cardiovascular Genetics* is online at:  
<http://circgenetics.ahajournals.org/subscriptions/>

## Supplemental Material.

**Norton et al, Circulation Cardiovascular Genetics 2013: *Exome sequencing and genome-wide linkage analysis in 17 families illustrates the complex contribution of TTN truncating variants to dilated cardiomyopathy.***

## Further acknowledgements

### HeartGO:

**Atherosclerosis Risk in Communities (ARIC):** NHLBI (N01 HC-55015, N01 HC-55016, N01HC-55017, N01 HC-55018, N01 HC-55019, N01 HC-55020, N01 HC-55021); **Cardiovascular Health Study (CHS):** NHLBI (N01-HC-85239, N01-HC-85079 through N01-HC-85086, N01-HC-35129, N01 HC-15103, N01 HC-55222, N01-HC-75150, N01-HC-45133, and grant HL080295), with additional support from NINDS and from NIA (AG-023629, AG-15928, AG-20098, and AG-027058); **Coronary Artery Risk Development in Young Adults (CARDIA):** NHLBI (N01-HC95095 & N01-HC48047, N01-HC48048, N01-HC48049, and N01-HC48050); **Framingham Heart Study (FHS):** NHLBI (N01-HC-25195 and grant R01 NS17950) with additional support from NIA (AG08122 and AG033193); **Jackson Heart Study (JHS):** NHLBI and the National Institute on Minority Health and Health Disparities (N01 HC-95170, N01 HC-95171 and N01 HC-95172); **Multi-Ethnic Study of Atherosclerosis (MESA):** NHLBI (N01-HC-95159 through N01-HC-95169 and RR-024156).

### Lung GO:

**Cystic Fibrosis (CF):** Cystic Fibrosis Foundation (GIBSON07K0, KNOWLE00A0, OBSERV04K0, RDP R026), the NHLBI (R01 HL-068890, R02 HL-095396), NIH National Center for Research Resources (UL1 RR-025014), and the National Human Genome Research Institute (NHGRI) (5R00 HG-004316). **Chronic Obstructive Pulmonary Disease (COPDGene):** NHLBI (U01 HL-089897, U01 HL-089856), and the COPD Foundation through contributions made to an Industry Advisory Board comprised of AstraZeneca, Boehringer Ingelheim, Novartis, Pfizer, and Sunovian. The COPDGene clinical centers and investigators are available at [www.copdgene.org](http://www.copdgene.org). **Acute Lung Injury (ALI):** NHLBI (RC2 HL-101779). **Lung Health Study (LHS):** NHLBI (RC2 HL-066583), the NHGRI (HG-004738), and the NHLBI Division of Lung Diseases (HR-46002). **Pulmonary Arterial Hypertension (PAH):** NIH (P50 HL-084946, K23 AR-52742), and the NHLBI (F32 HL-083714). **Asthma:** NHLBI (RC2 HL-101651), and the NIH (HL-077916, HL-69197, HL-76285, M01 RR-07122).

### SWISS and ISGS:

Siblings with Ischemic Stroke Study (SWISS): National Institute of Neurological Disorders and Stroke (NINDS) (R01 NS039987); Ischemic Stroke Genetics Study (ISGS): NINDS (R01 NS042733)

### WHISP:



**Women's Health Initiative (WHI):** The WHI Sequencing Project is funded by the NHLBI (HL-102924) as well as the National Institutes of Health (NIH), U.S. Department of Health and Human Services through contracts N01WH22110, 24152, 32100-2, 32105-6, 32108-9, 32111-13, 32115, 32118-32119, 32122, 42107-26, 42129-32, and 44221. The authors thank the WHI investigators and staff for their dedication, and the study participants for making the program possible. A full listing of WHI investigators can be found at: [http://www.whiscience.org/publications/WHI\\_investigators\\_shortlist.pdf](http://www.whiscience.org/publications/WHI_investigators_shortlist.pdf)

## NHLBI GO Exome Sequencing Project

### BroadGO

Stacey B. Gabriel (Broad Institute)<sup>4, 5, 11, 16, 17</sup>, David M. Altshuler (Broad Institute, Harvard Medical School, Massachusetts General Hospital)<sup>1, 5, 7, 17</sup>, Gonçalo R. Abecasis (University of Michigan)<sup>3, 5, 9, 13, 15, 17</sup>, Hooman Allayee (University of Southern California)<sup>5</sup>, Sharon Cresci (Washington University School of Medicine)<sup>5</sup>, Mark J. Daly (Broad Institute, Massachusetts General Hospital), Paul I. W. de Bakker (Broad Institute, Harvard Medical School, University Medical Center Utrecht)<sup>3, 15</sup>, Mark A. DePristo (Broad Institute)<sup>4, 13, 15, 16</sup>, Ron Do (Broad Institute)<sup>5, 9, 13, 15</sup>, Peter Donnelly (University of Oxford)<sup>5</sup>, Deborah N. Farlow (Broad Institute)<sup>3, 4, 5, 12, 14, 16, 17</sup>, Tim Fennell (Broad Institute), Kiran Garimella (University of Oxford)<sup>4, 16</sup>, Stanley L. Hazen (Cleveland Clinic)<sup>5</sup>, Youna Hu (University of Michigan)<sup>3, 9, 15</sup>, Daniel M. Jordan (Harvard Medical School, Harvard University)<sup>13</sup>, Goo Jun (University of Michigan)<sup>13</sup>, Sekar Kathiresan (Broad Institute, Harvard Medical School, Massachusetts General Hospital)<sup>5, 8, 9, 12, 14, 15, 17, 20</sup>, Hyun Min Kang (University of Michigan)<sup>9, 13, 16</sup>, Adam Kiezun (Broad Institute)<sup>5, 13, 15</sup>, Guillaume Lettre (Broad Institute, Montreal Heart Institute, Université de Montréal)<sup>1, 2, 13, 15</sup>, Bingshan Li (University of Michigan)<sup>3</sup>, Mingyao Li (University of Pennsylvania)<sup>5</sup>, Christopher H. Newton-Cheh (Broad Institute, Massachusetts General Hospital, Harvard Medical School)<sup>3, 8, 15</sup>, Sandosh Padmanabhan (University of Glasgow School of Medicine)<sup>3, 12, 15</sup>, Gina Peloso (Broad Institute, Harvard Medical School, Massachusetts General Hospital)<sup>5</sup>, Sara Pulit (Broad Institute)<sup>3, 15</sup>, Daniel J. Rader (University of Pennsylvania)<sup>5</sup>, David Reich (Broad Institute, Harvard Medical School)<sup>15</sup>, Muredach P. Reilly (University of Pennsylvania)<sup>5</sup>, Manuel A. Rivas (Broad Institute, Massachusetts General Hospital)<sup>5</sup>, Steve Schwartz (Fred Hutchinson Cancer Research Center)<sup>5, 12</sup>, Laura Scott (University of Michigan)<sup>1</sup>, David S. Siscovick (University of Washington)<sup>5, 1, 25</sup>, John A. Spertus (University of Missouri Kansas City)<sup>5</sup>, Nathaniel O. Stitzel (Brigham and Women's Hospital)<sup>5, 15</sup>, Nina Stoletzki (Brigham and Women's Hospital, Broad Institute, Harvard Medical School)<sup>13</sup>, Shamil R. Sunyaev (Brigham and Women's Hospital, Broad Institute, Harvard Medical School)<sup>1, 3, 5, 13, 15</sup>, Benjamin F. Voight (Broad Institute, Massachusetts General Hospital), Cristen J. Willer (University of Michigan)<sup>1, 9, 13, 15</sup>

### HeartGO

Stephen S. Rich (University of Virginia)<sup>2, 4, 7, 8, 9, 11, 14, 15, 17, 18, 31</sup>, Ermeg Akylbekova (Jackson State University, University of Mississippi Medical Center)<sup>29</sup>, Larry D. Atwood\* (Boston University)<sup>1, 11, 28</sup>, Christie M. Ballantyne (Baylor College of Medicine, Methodist DeBakey Heart Center)<sup>9, 22</sup>, Maja Barbalic (University of Texas Health Science Center Houston)<sup>9, 14, 15, 17, 22</sup>, R. Graham Barr (Columbia University Medical Center)<sup>10, 31</sup>, Emelia J. Benjamin (Boston University)<sup>14, 20, 28</sup>, Joshua Bis (University of Washington)<sup>15, 23</sup>, Eric Boerwinkle (University of Texas Health Science Center Houston)<sup>3, 5, 9, 13, 15, 17, 22</sup>, Donald

W. Bowden (Wake Forest University)<sup>1, 31</sup>, Jennifer Brody (University of Washington)<sup>3, 5, 15, 23</sup>, Matthew Budoff (Harbor-UCLA Medical Center)<sup>31</sup>, Greg Burke (Wake Forest University)<sup>5, 31</sup>, Sarah Buxbaum (Jackson State University)<sup>3, 13, 15, 29</sup>, Jeff Carr (Wake Forest University)<sup>25, 29, 31</sup>, Donna T. Chen (University of Virginia)<sup>6, 11</sup>, Ida Y. Chen (Cedars-Sinai Medical Center)<sup>1, 31</sup>, Wei-Min Chen (University of Virginia)<sup>13, 15, 18</sup>, Pat Concannon (University of Virginia)<sup>11</sup>, Jacy Crosby (University of Texas Health Science Center Houston)<sup>22</sup>, L. Adrienne Cupples (Boston University)<sup>1, 3, 5, 9, 13, 15, 18, 28</sup>, Ralph D'Agostino (Boston University)<sup>28</sup>, Anita L. DeStefano (Boston University)<sup>13, 18, 28</sup>, Albert Dreisbach (University of Mississippi Medical Center)<sup>3, 29</sup>, Josée Dupuis (Boston University)<sup>1, 28</sup>, J. Peter Durda (University of Vermont)<sup>15, 23</sup>, Jaclyn Ellis (University of North Carolina Chapel Hill)<sup>1</sup>, Aaron R. Folsom (University of Minnesota)<sup>5, 22</sup>, Myriam Fornage (University of Texas Health Science Center Houston)<sup>3, 18, 25</sup>, Caroline S. Fox (National Heart, Lung, and Blood Institute)<sup>1, 28</sup>, Ervin Fox (University of Mississippi Medical Center)<sup>3, 9, 29</sup>, Vincent Funari (Cedars-Sinai Medical Center)<sup>1, 11, 31</sup>, Santhi K. Ganesh (University of Michigan)<sup>2, 22</sup>, Julius Gardin (Hackensack University Medical Center)<sup>25</sup>, David Goff (Wake Forest University)<sup>25</sup>, Ora Gordon (Cedars-Sinai Medical Center)<sup>11, 31</sup>, Wayne Grody (University of California Los Angeles)<sup>11, 31</sup>, Myron Gross (University of Minnesota)<sup>1, 5, 14, 25</sup>, Xiuqing Guo (Cedars-Sinai Medical Center)<sup>3, 15, 31</sup>, Ira M. Hall (University of Virginia), Nancy L. Heard-Costa (Boston University)<sup>1, 11, 28</sup>, Susan R. Heckbert (University of Washington)<sup>10, 14, 20, 23</sup>, Nicholas Heintz (University of Vermont), David M. Herrington (Wake Forest University)<sup>5, 31</sup>, DeMarc Hickson (Jackson State University, University of Mississippi Medical Center)<sup>29</sup>, Jie Huang (National Heart, Lung, and Blood Institute)<sup>5, 28</sup>, Shih-Jen Hwang (Boston University, National Heart, Lung, and Blood Institute)<sup>3, 28</sup>, David R. Jacobs (University of Minnesota)<sup>25</sup>, Nancy S. Jenny (University of Vermont)<sup>1, 2, 23</sup>, Andrew D. Johnson (National Heart, Lung, and Blood Institute)<sup>2, 5, 11, 28</sup>, Craig W. Johnson (University of Washington)<sup>15, 31</sup>, Steven Kawut (University of Pennsylvania)<sup>10, 31</sup>, Richard Kronmal (University of Washington)<sup>31</sup>, Raluca Kurz (Cedars-Sinai Medical Center)<sup>11, 31</sup>, Ethan M. Lange (University of North Carolina Chapel Hill)<sup>3, 5, 9, 13, 34</sup>, Leslie A. Lange (University of North Carolina Chapel Hill)<sup>1, 2, 3, 5, 9, 12, 13, 15, 17, 18, 20, 25, 34</sup>, Martin G. Larson (Boston University)<sup>3, 15, 28</sup>, Mark Lawson (University of Virginia), Cora E. Lewis (University of Alabama at Birmingham)<sup>25, 34</sup>, Daniel Levy (National Heart, Lung, and Blood Institute)<sup>3, 15, 17, 28</sup>, Dalin Li (Cedars-Sinai Medical Center)<sup>11, 15, 31</sup>, Honghuang Lin (Boston University)<sup>20, 28</sup>, Chunyu Liu (National Heart, Lung, and Blood Institute)<sup>3, 28</sup>, Jiankang Liu (University of Mississippi Medical Center)<sup>1, 29</sup>, Kiang Liu (Northwestern University)<sup>25</sup>, Xiaoming Liu (University of Texas Health Science Center Houston)<sup>15, 22</sup>, Yongmei Liu (Wake Forest University)<sup>2, 5, 31</sup>, William T. Longstreth (University of Washington)<sup>18, 23</sup>, Cay Loria (National Heart, Lung, and Blood Institute)<sup>25</sup>, Thomas Lumley (University of Auckland)<sup>9, 23</sup>, Kathryn Lunetta (Boston University)<sup>28</sup>, Aaron J. Mackey (University of Virginia)<sup>16, 18</sup>, Rachel Mackey (University of Pittsburgh)<sup>1, 23, 31</sup>, Ani Manichaikul (University of Virginia)<sup>8, 15, 18, 31</sup>, Taylor Maxwell (University of Texas Health Science Center Houston)<sup>22</sup>, Barbara McKnight (University of Washington)<sup>15, 23</sup>, James B. Meigs (Brigham and Women's Hospital, Harvard Medical School, Massachusetts General Hospital)<sup>1, 28</sup>, Alanna C. Morrison (University of Texas Health Science Center Houston)<sup>3, 15, 17</sup>, Solomon K. Musani (University of Mississippi Medical Center)<sup>3, 29</sup>, Josyf C. Mychaleckyj (University of Virginia)<sup>13, 15, 31</sup>, Jennifer A. Nettleton (University of Texas Health Science Center Houston)<sup>9, 22</sup>, Kari North (University of North Carolina Chapel Hill)<sup>1, 3, 9, 10, 13, 15, 17, 34</sup>, Christopher J. O'Donnell (Massachusetts General Hospital, National Heart, Lung, and Blood Institute)<sup>2, 5, 9, 11, 12, 14, 15, 17, 20, 28</sup>, Daniel O'Leary (Tufts University School of Medicine)<sup>25, 31</sup>, Frank Ong (Cedars-Sinai Medical Center)<sup>3, 11, 31</sup>, Walter Palmas (Columbia University)<sup>3, 15, 31</sup>, James S. Pankow (University of Minnesota)<sup>1, 22</sup>, Nathan D. Pankratz (Indiana University School of Medicine)<sup>15, 25</sup>, Shom

Paul (University of Virginia), Marco Perez (Stanford University School of Medicine), Sharina D. Person (University of Alabama at Birmingham, University of Alabama at Tuscaloosa)<sup>25</sup>, Joseph Polak (Tufts University School of Medicine)<sup>31</sup>, Wendy S. Post (Johns Hopkins University)<sup>3, 9, 11, 14, 20, 31</sup>, Bruce M. Psaty (Group Health Research Institute, University of Washington)<sup>3, 5, 9, 11, 14, 15, 23</sup>, Aaron R. Quinlan (University of Virginia)<sup>18, 19</sup>, Leslie J. Raffel (Cedars-Sinai Medical Center)<sup>6, 11, 31</sup>, Vasam S. Ramachandran (Boston University)<sup>3, 28</sup>, Alexander P. Reiner (Fred Hutchinson Cancer Research Center, University of Washington)<sup>1, 2, 3, 5, 9, 11, 12, 13, 14, 15, 20, 25, 34</sup>, Kenneth Rice (University of Washington)<sup>15, 23</sup>, Jerome I. Rotter (Cedars-Sinai Medical Center)<sup>1, 3, 6, 8, 11, 15, 31</sup>, Jill P. Sanders (University of Vermont)<sup>23</sup>, Pamela Schreiner (University of Minnesota)<sup>25</sup>, Sudha Seshadri (Boston University)<sup>18, 28</sup>, Steve Shea (Brigham and Women's Hospital, Harvard University)<sup>28</sup>, Stephen Sidney (Kaiser Permanente Division of Research, Oakland, CA)<sup>25</sup>, Kevin Silverstein (University of Minnesota)<sup>25</sup>, David S. Siscovick (University of Washington)<sup>5, 1, 25</sup>, Nicholas L. Smith (University of Washington)<sup>2, 15, 20, 23</sup>, Nona Sotoodehnia (University of Washington)<sup>3, 15, 23</sup>, Asoke Srinivasan (Tougaloo College)<sup>29</sup>, Herman A. Taylor (Jackson State University, Tougaloo College, University of Mississippi Medical Center)<sup>5, 29</sup>, Kent Taylor (Cedars-Sinai Medical Center)<sup>31</sup>, Fridtjof Thomas (University of Texas Health Science Center Houston)<sup>3, 22</sup>, Russell P. Tracy (University of Vermont)<sup>5, 9, 11, 12, 14, 15, 17, 20, 23</sup>, Michael Y. Tsai (University of Minnesota)<sup>9, 31</sup>, Kelly A. Volcik (University of Texas Health Science Center Houston)<sup>22</sup>, Christina L. Wassel (University of California San Diego)<sup>9, 15, 31</sup>, Karol Watson (University of California Los Angeles)<sup>31</sup>, Gina Wei (National Heart, Lung, and Blood Institute)<sup>25</sup>, Wendy White (Tougaloo College)<sup>29</sup>, Kerri L. Wiggins (University of Vermont)<sup>23</sup>, Jemma B. Wilk (Boston University)<sup>10, 28</sup>, O. Dale Williams (Florida International University)<sup>25</sup>, Gregory Wilson (Jackson State University)<sup>29</sup>, James G. Wilson (University of Mississippi Medical Center)<sup>1, 2, 5, 8, 9, 11, 12, 14, 17, 20, 29</sup>, Phillip Wolf (Boston University)<sup>28</sup>, Neil A. Zakai (University of Vermont)<sup>2, 23</sup>

### **ISGS and SWISS**

John Hardy (Reta Lila Weston Research Laboratories, Institute of Neurology, University College London)<sup>18</sup>, James F. Meschia (Mayo Clinic)<sup>18</sup>, Michael Nalls (National Institute on Aging)<sup>2, 18</sup>, Stephen S. Rich (University of Virginia)<sup>2, 4, 7, 8, 9, 11, 14, 15, 17, 18, 31</sup>, Andrew Singleton (National Institute on Aging)<sup>18</sup>, Brad Worrall (University of Virginia)<sup>18</sup>

### **LungGO**

Michael J. Bamshad (Seattle Children's Hospital, University of Washington)<sup>4, 6, 7, 8, 10, 11, 13, 15, 17, 27</sup>, Kathleen C. Barnes (Johns Hopkins University)<sup>2, 10, 12, 14, 15, 17, 20, 24, 30, 32</sup>, Ibrahim Abdulhamid (Children's Hospital of Michigan)<sup>27</sup>, Frank Accurso (University of Colorado)<sup>27</sup>, Ran Anbar (Upstate Medical University)<sup>27</sup>, Terri Beaty (Johns Hopkins University)<sup>24, 30</sup>, Abigail Bigham (University of Washington)<sup>13, 15, 27</sup>, Phillip Black (Children's Mercy Hospital)<sup>27</sup>, Eugene Bleeker (Wake Forest University)<sup>33</sup>, Kati Buckingham (University of Washington)<sup>27</sup>, Anne Marie Cairns (Maine Medical Center)<sup>27</sup>, Wei-Min Chen (University of Virginia)<sup>13, 15, 18</sup>, Daniel Caplan (Emory University)<sup>27</sup>, Barbara Chatfield (University of Utah)<sup>27</sup>, Aaron Chidekel (A.I. Dupont Institute Medical Center)<sup>27</sup>, Michael Cho (Brigham and Women's Hospital, Harvard Medical School)<sup>13, 15, 24</sup>, David C. Christiani (Massachusetts General Hospital)<sup>21</sup>, James D. Crapo (National Jewish Health)<sup>24, 30</sup>, Julia Crouch (Seattle Children's Hospital)<sup>6</sup>, Denise Daley (University of British Columbia)<sup>30</sup>, Anthony Dang (University of North Carolina Chapel Hill)<sup>26</sup>, Hong Dang (University of North Carolina Chapel Hill)<sup>26</sup>, Alicia De Paula (Ochsner Health System)<sup>27</sup>, Joan DeCelie-Germana (Schneider Children's Hospital)<sup>27</sup>, Allen Dozor (New York Medical College, Westchester Medical Center)<sup>27</sup>, Mitch Drumm (University of North

Carolina Chapel Hill)<sup>26</sup>, Maynard Dyson (Cook Children's Med. Center)<sup>27</sup>, Julia Emerson (Seattle Children's Hospital, University of Washington)<sup>27</sup>, Mary J. Emond (University of Washington)<sup>10, 13, 15, 17, 27</sup>, Thomas Ferkol (St. Louis Children's Hospital, Washington University School of Medicine)<sup>27</sup>, Robert Fink (Children's Medical Center of Dayton)<sup>27</sup>, Cassandra Foster (Johns Hopkins University)<sup>30</sup>, Deborah Froh (University of Virginia)<sup>27</sup>, Li Gao (Johns Hopkins University)<sup>24, 30, 32</sup>, William Gershan (Children's Hospital of Wisconsin)<sup>27</sup>, Ronald L. Gibson (Seattle Children's Hospital, University of Washington)<sup>10, 27</sup>, Elizabeth Godwin (University of North Carolina Chapel Hill)<sup>26</sup>, Magdalen Gondor (All Children's Hospital Cystic Fibrosis Center)<sup>27</sup>, Hector Gutierrez (University of Alabama at Birmingham)<sup>27</sup>, Nadia N. Hansel (Johns Hopkins University, Johns Hopkins University School of Public Health)<sup>10, 15, 30</sup>, Paul M. Hassoun (Johns Hopkins University)<sup>10, 14, 32</sup>, Peter Hiatt (Texas Children's Hospital)<sup>27</sup>, John E. Hokanson (University of Colorado)<sup>24</sup>, Michelle Howenstine (Indiana University, Riley Hospital for Children)<sup>27</sup>, Laura K. Hummer (Johns Hopkins University)<sup>32</sup>, Jamshed Kanga (University of Kentucky)<sup>27</sup>, Yoonhee Kim (National Human Genome Research Institute)<sup>24, 32</sup>, Michael R. Knowles (University of North Carolina Chapel Hill)<sup>10, 26</sup>, Michael Konstan (Rainbow Babies & Children's Hospital)<sup>27</sup>, Thomas Lahiri (Vermont Children's Hospital at Fletcher Allen Health Care)<sup>27</sup>, Nan Laird (Harvard School of Public Health)<sup>24</sup>, Christoph Lange (Harvard School of Public Health)<sup>24</sup>, Lin Lin (Harvard Medical School)<sup>21</sup>, Xihong Lin (Harvard School of Public Health)<sup>21</sup>, Tin L. Louie (University of Washington)<sup>13, 15, 27</sup>, David Lynch (National Jewish Health)<sup>24</sup>, Barry Make (National Jewish Health)<sup>24</sup>, Thomas R. Martin (University of Washington, VA Puget Sound Medical Center)<sup>10, 21</sup>, Steve C. Mathai (Johns Hopkins University)<sup>32</sup>, Rasika A. Mathias (Johns Hopkins University)<sup>10, 13, 15, 30, 32</sup>, John McNamara (Children's Hospitals and Clinics of Minnesota)<sup>27</sup>, Sharon McNamara (Seattle Children's Hospital)<sup>27</sup>, Deborah Meyers (Wake Forest University)<sup>33</sup>, Susan Millard (DeVos Children's Butterworth Hospital, Spectrum Health Systems)<sup>27</sup>, Peter Mogayzel (Johns Hopkins University)<sup>27</sup>, Richard Moss (Stanford University)<sup>27</sup>, Tanda Murray (Johns Hopkins University)<sup>30</sup>, Dennis Nielson (University of California at San Francisco)<sup>27</sup>, Blakeslee Noyes (Cardinal Glennon Children's Hospital)<sup>27</sup>, Wanda O'Neal (University of North Carolina Chapel Hill)<sup>26</sup>, David Orenstein (Children's Hospital of Pittsburgh)<sup>27</sup>, Brian O'Sullivan (University of Massachusetts Memorial Health Care)<sup>27</sup>, Rhonda Pace (University of North Carolina Chapel Hill)<sup>26</sup>, Peter Pare (St. Paul's Hospital)<sup>30</sup>, H. Worth Parker (Dartmouth-Hitchcock Medical Center, New Hampshire Cystic Fibrosis Center)<sup>27</sup>, Mary Ann Passero (Rhode Island Hospital)<sup>27</sup>, Elizabeth Perket (Vanderbilt University)<sup>27</sup>, Adrienne Prestridge (Children's Memorial Hospital)<sup>27</sup>, Nicholas M. Rafaels (Johns Hopkins University)<sup>30</sup>, Bonnie Ramsey (Seattle Children's Hospital, University of Washington)<sup>27</sup>, Elizabeth Regan (National Jewish Health)<sup>24</sup>, Clement Ren (University of Rochester)<sup>27</sup>, George Retsch-Bogart (University of North Carolina Chapel Hill)<sup>27</sup>, Michael Rock (University of Wisconsin Hospital and Clinics)<sup>27</sup>, Antony Rosen (Johns Hopkins University)<sup>32</sup>, Margaret Rosenfeld (Seattle Children's Hospital, University of Washington)<sup>27</sup>, Ingo Ruczinski (Johns Hopkins University School of Public Health)<sup>13, 15, 30</sup>, Andrew Sanford (University of British Columbia)<sup>30</sup>, David Schaeffer (Nemours Children's Clinic)<sup>27</sup>, Cindy Sell (University of North Carolina Chapel Hill)<sup>26</sup>, Daniel Sheehan (Children's Hospital of Buffalo)<sup>27</sup>, Edwin K. Silverman (Brigham and Women's Hospital, Harvard Medical School)<sup>24, 30</sup>, Don Sin (Children's Medical Center of Dayton)<sup>30</sup>, Terry Spencer (Elliot Health System)<sup>27</sup>, Jackie Stonebraker (University of North Carolina Chapel Hill)<sup>26</sup>, Holly K. Tabor (Seattle Children's Hospital, University of Washington)<sup>6, 10, 11, 17, 27</sup>, Laurie Varlotta (St. Christopher's Hospital for Children)<sup>27</sup>, Candelaria I. Vergara (Johns Hopkins University)<sup>30</sup>, Robert Weiss<sup>30</sup>, Fred Wigley (Johns Hopkins University)<sup>32</sup>, Robert A. Wise (Johns Hopkins University)<sup>30</sup>, Fred A. Wright (University of North



Carolina Chapel Hill)<sup>26</sup>, Mark M. Wurfel (University of Washington)<sup>10, 14, 21</sup>, Robert Zanni (Monmouth Medical Center)<sup>27</sup>, Fei Zou (University of North Carolina Chapel Hill)<sup>26</sup>

### SeattleGO

Deborah A. Nickerson (University of Washington)<sup>3, 4, 5, 7, 8, 9, 11, 15, 17, 18, 19</sup>, Mark J. Rieder (University of Washington)<sup>4, 11, 13, 15, 16, 17, 19</sup>, Phil Green (University of Washington), Jay Shendure (University of Washington)<sup>1, 8, 14, 16, 17</sup>, Joshua M. Akey (University of Washington)<sup>13, 14, 15</sup>, Michael J. Bamshad (Seattle Children's Hospital, University of Washington)<sup>4, 6, 7, 8, 10, 11, 13, 15, 17, 27</sup>, Carlos D. Bustamante (Stanford University School of Medicine)<sup>3, 13, 15</sup>, David R. Crosslin (University of Washington)<sup>2, 9</sup>, Evan E. Eichler (University of Washington)<sup>19</sup>, P. Keolu Fox<sup>2</sup>, Wenqing Fu (University of Washington)<sup>13</sup>, Adam Gordon (University of Washington)<sup>11</sup>, Simon Gravel (Stanford University School of Medicine)<sup>13, 15</sup>, Gail P. Jarvik (University of Washington)<sup>9, 15</sup>, Jill M. Johnsen (Puget Sound Blood Center, University of Washington)<sup>2</sup>, Mengyuan Kan (Baylor College of Medicine)<sup>13</sup>, Eimear E. Kenny (Stanford University School of Medicine)<sup>3, 13, 15</sup>, Jeffrey M. Kidd (Stanford University School of Medicine)<sup>13, 15</sup>, Fremiet Lara-Garduno (Baylor College of Medicine)<sup>15</sup>, Suzanne M. Leal (Baylor College of Medicine)<sup>1, 13, 15, 16, 17, 19, 20</sup>, Dajiang J. Liu (Baylor College of Medicine)<sup>13, 15</sup>, Sean McGee (University of Washington)<sup>13, 15, 19</sup>, Timothy D. O'Connor (University of Washington)<sup>13</sup>, Bryan Paepers (University of Washington)<sup>16</sup>, Peggy D. Robertson (University of Washington)<sup>4</sup>, Joshua D. Smith (University of Washington)<sup>4, 16, 19</sup>, Jeffrey C. Staples (University of Washington), Jacob A. Tennessen (University of Washington)<sup>13</sup>, Emily H. Turner (University of Washington)<sup>4, 16</sup>, Gao Wang (Baylor College of Medicine)<sup>1, 13, 20</sup>, Qian Yi (University of Washington)<sup>4</sup>

### WHISP

Rebecca Jackson (Ohio State University)<sup>1, 2, 4, 5, 8, 12, 14, 15, 17, 18, 20, 34</sup>, Kari North (University of North Carolina Chapel Hill)<sup>1, 3, 9, 10, 13, 15, 17, 34</sup>, Ulrike Peters (Fred Hutchinson Cancer Research Center)<sup>1, 3, 11, 12, 13, 15, 17, 18, 34</sup>, Christopher S. Carlson (Fred Hutchinson Cancer Research Center, University of Washington)<sup>1, 2, 3, 5, 12, 13, 14, 15, 16, 17, 18, 19, 34</sup>, Garnet Anderson (Fred Hutchinson Cancer Research Center)<sup>34</sup>, Hoda Anton-Culver (University of California at Irvine)<sup>34</sup>, Themistocles L. Assimes (Stanford University School of Medicine)<sup>5, 9, 11, 34</sup>, Paul L. Auer (Fred Hutchinson Cancer Research Center)<sup>1, 2, 3, 5, 11, 12, 13, 15, 16, 18, 34</sup>, Shirley Beresford (Fred Hutchinson Cancer Research Center)<sup>34</sup>, Chris Bizon (University of North Carolina Chapel Hill)<sup>3, 9, 13, 15, 34</sup>, Henry Black (Rush Medical Center)<sup>34</sup>, Robert Brunner (University of Nevada)<sup>34</sup>, Robert Brzyski (University of Texas Health Science Center San Antonio)<sup>34</sup>, Dale Burwen (National Heart, Lung, and Blood Institute WHI Project Office)<sup>34</sup>, Bette Caan (Kaiser Permanente Division of Research, Oakland, CA)<sup>34</sup>, Cara L. Carty (Fred Hutchinson Cancer Research Center)<sup>18, 34</sup>, Rowan Chlebowski (Los Angeles Biomedical Research Institute)<sup>34</sup>, Steven Cummings (University of California at San Francisco)<sup>34</sup>, J. David Curb\* (University of Hawaii)<sup>9, 18, 34</sup>, Charles B. Eaton (Brown University, Memorial Hospital of Rhode Island)<sup>12, 34</sup>, Leslie Ford (National Heart, Lung, and Blood Institute, National Heart, Lung, and Blood Institute WHI Project Office)<sup>34</sup>, Nora Franceschini (University of North Carolina Chapel Hill)<sup>2, 3, 9, 10, 15, 34</sup>, Stephanie M. Fullerton (University of Washington)<sup>6, 11, 34</sup>, Margery Gass (University of Cincinnati)<sup>34</sup>, Nancy Geller (National Heart, Lung, and Blood Institute WHI Project Office)<sup>34</sup>, Gerardo Heiss (University of North Carolina Chapel Hill)<sup>5, 34</sup>, Barbara V. Howard (Howard University, MedStar Research Institute)<sup>34</sup>, Li Hsu (Fred Hutchinson Cancer Research Center)<sup>1, 13, 15, 18, 34</sup>, Carolyn M. Hutter (Fred Hutchinson Cancer

Research Center)<sup>13, 15, 18, 34</sup>, John Ioannidis (Stanford University School of Medicine)<sup>11, 34</sup>, Shuo Jiao (Fred Hutchinson Cancer Research Center)<sup>34</sup>, Karen C. Johnson (University of Tennessee Health Science Center)<sup>3, 34</sup>, Charles Kooperberg (Fred Hutchinson Cancer Research Center)<sup>1, 5, 9, 13, 14, 15, 17, 18, 34</sup>, Lewis Kuller (University of Pittsburgh)<sup>34</sup>, Andrea LaCroix (Fred Hutchinson Cancer Research Center)<sup>34</sup>, Kamakshi Lakshminarayan (University of Minnesota)<sup>18, 34</sup>, Dorothy Lane (State University of New York at Stony Brook)<sup>34</sup>, Ethan M. Lange (University of North Carolina Chapel Hill)<sup>3, 5, 9, 13, 34</sup>, Leslie A. Lange (University of North Carolina Chapel Hill)<sup>1, 2, 3, 5, 9, 12, 13, 15, 17, 18, 20, 25, 34</sup>, Norman Lasser (University of Medicine and Dentistry of New Jersey)<sup>34</sup>, Erin LeBlanc (Kaiser Permanente Center for Health Research, Portland, OR)<sup>34</sup>, Cora E. Lewis (University of Alabama at Birmingham)<sup>25, 34</sup>, Kuo-Ping Li (University of North Carolina Chapel Hill)<sup>9, 34</sup>, Marian Limacher (University of Florida)<sup>34</sup>, Dan-Yu Lin (University of North Carolina Chapel Hill)<sup>1, 3, 9, 13, 15, 34</sup>, Benjamin A. Logsdon (Fred Hutchinson Cancer Research Center)<sup>2, 34</sup>, Shari Ludlam (National Heart, Lung, and Blood Institute WHI Project Office)<sup>34</sup>, JoAnn E. Manson (Brigham and Women's Hospital, Harvard School of Public Health)<sup>34</sup>, Karen Margolis (University of Minnesota)<sup>34</sup>, Lisa Martin (George Washington University Medical Center)<sup>9, 34</sup>, Joan McGowan (National Heart, Lung, and Blood Institute WHI Project Office)<sup>34</sup>, Keri L. Monda (Amgen, Inc.)<sup>1, 15, 34</sup>, Jane Morley Kotchen (Medical College of Wisconsin)<sup>34</sup>, Lauren Nathan (University of California Los Angeles)<sup>34</sup>, Judith Ockene (Fallon Clinic, University of Massachusetts)<sup>34</sup>, Mary Jo O'Sullivan (University of Miami)<sup>34</sup>, Lawrence S. Phillips (Emory University)<sup>34</sup>, Ross L. Prentice (Fred Hutchinson Cancer Research Center)<sup>34</sup>, Alexander P. Reiner (Fred Hutchinson Cancer Research Center, University of Washington)<sup>1, 2, 3, 5, 9, 11, 12, 13, 14, 15, 20, 25, 34</sup>, John Robbins (University of California at Davis)<sup>34</sup>, Jennifer G. Robinson (University of Iowa)<sup>9, 11, 18, 34</sup>, Jacques E. Rossouw (National Heart, Lung, and Blood Institute, National Heart, Lung, and Blood Institute WHI Project Office)<sup>5, 14, 17, 20, 34</sup>, Haleh Sangi-Haghpeykar (Baylor College of Medicine)<sup>34</sup>, Gloria E. Sarto (University of Wisconsin)<sup>34</sup>, Sally Shumaker (Wake Forest University)<sup>34</sup>, Michael S. Simon (Wayne State University)<sup>34</sup>, Marcia L. Stefanick (Stanford University School of Medicine)<sup>34</sup>, Evan Stein (Medical Research Labs)<sup>34</sup>, Hua Tang (Stanford University)<sup>2, 34</sup>, Kira C. Taylor (University of Louisville)<sup>1, 3, 13, 15, 20, 34</sup>, Cynthia A. Thomson (University of Arizona)<sup>34</sup>, Timothy A. Thornton (University of Washington)<sup>13, 15, 18, 34</sup>, Linda Van Horn (Northwestern University)<sup>34</sup>, Mara Vitolins (Wake Forest University)<sup>34</sup>, Jean Wactawski-Wende (University of Buffalo)<sup>34</sup>, Robert Wallace (University of Iowa)<sup>2, 34</sup>, Sylvia Wassertheil-Smoller (Boston University)<sup>18, 34</sup>, Donglin Zeng (University of North Carolina Chapel Hill)<sup>9, 34</sup>

#### **NHLBI GO ESP Project Team**

Deborah Applebaum-Bowden (National Heart, Lung, and Blood Institute)<sup>4, 7, 12, 17</sup>, Michael Feolo (National Center for Biotechnology Information)<sup>12</sup>, Weiniu Gan (National Heart, Lung, and Blood Institute)<sup>7, 8, 16, 17</sup>, Dina N. Paltoo (National Heart, Lung, and Blood Institute)<sup>4, 6, 11, 17</sup>, Jacques E. Rossouw (National Heart, Lung, and Blood Institute, National Heart, Lung, and Blood Institute WHI Project Office)<sup>5, 14, 17, 20, 34</sup>, Phyllis Sholinsky (National Heart, Lung, and Blood Institute)<sup>4, 12, 17</sup>, Anne Sturcke (National Center for Biotechnology Information)<sup>12</sup>

\*deceased

#### **ESP Groups**

<sup>1</sup>Anthropometry Project Team, <sup>2</sup>Blood Count/Hematology Project Team, <sup>3</sup>Blood Pressure Project Team, <sup>4</sup>Data Flow Working Group, <sup>5</sup>Early MI Project Team, <sup>6</sup>ELSI Working Group, <sup>7</sup>Executive Committee, <sup>8</sup>Family Study Project Team, <sup>9</sup>Lipids Project Team,

<sup>10</sup>Lung Project Team, <sup>11</sup>Personal Genomics Project Team, <sup>12</sup>Phenotype and Harmonization Working Group, <sup>13</sup>Population Genetics and Statistical Analysis Working Group, <sup>14</sup>Publications and Presentations Working Group, <sup>15</sup>Quantitative Analysis Ad Hoc Task Group, <sup>16</sup>Sequencing and Genotyping Working Group, <sup>17</sup>Steering Committee, <sup>18</sup>Stroke Project Team, <sup>19</sup>Structural Variation Working Group, <sup>20</sup>Subclinical/Quantitative Project Team

**ESP Cohorts**

<sup>21</sup>Acute Lung Injury (ALI), <sup>22</sup>Atherosclerosis Risk in Communities (ARIC), <sup>23</sup>Cardiovascular Health Study (CHS), <sup>24</sup>Chronic Obstructive Pulmonary Disease (COPDGene), <sup>25</sup>Coronary Artery Risk Development in Young Adults (CARDIA), <sup>26</sup>Cystic Fibrosis (CF), <sup>27</sup>Early Pseudomonas Infection Control (EPIC), <sup>28</sup>Framingham Heart Study (FHS), <sup>29</sup>Jackson Heart Study (JHS), <sup>30</sup>Lung Health Study (LHS), <sup>31</sup>Multi-Ethnic Study of Atherosclerosis (MESA), <sup>32</sup>Pulmonary Arterial Hypertension (PAH), <sup>33</sup>Severe Asthma Research Program (SARP), <sup>34</sup>Women's Health Initiative (WHI)