

## New Insights Into the Genetic Basis of Inherited Arrhythmia Syndromes

Belinda Gray, BSc (Med), MBBS; Elijah R. Behr, MA, MBBS, MD

Inherited arrhythmia syndromes encompass several different diseases, including long QT syndrome (LQTS), Brugada syndrome (BrS), catecholaminergic polymorphic ventricular tachycardia (CPVT), short QT syndrome (SQTS), idiopathic ventricular fibrillation (IVF), and progressive cardiac conduction system disease (PCCD).<sup>1</sup> The heart is typically structurally normal with no evidence of disease macroscopically. They are an important cause for sudden cardiac death in the young, and an autopsy is typically negative.<sup>2,3</sup>

Ventricular arrhythmias are caused by mutations of ion channels and their interacting proteins, predominantly involving potassium, sodium, and calcium handling.<sup>4</sup> Genetic studies have identified the specific genetic abnormalities that underpin these diseases, even permitting diagnosis in the deceased using post-mortem genetic testing (the molecular autopsy).<sup>3</sup> Most arrhythmia syndromes are inherited in an autosomal dominant manner, such that first-degree family members have a 50% chance of inheriting the disease. Identification of the mutation allows for predictive genetic testing in other living family members.<sup>4</sup> Variable penetrance is common in all arrhythmia syndromes, the same mutation in the same family causing wide variation in phenotype.<sup>4</sup> This suggests that other factors such as genetic modifiers and environmental factors may influence the phenotype.

This review will highlight the latest developments in understanding the genetic basis of inherited arrhythmia syndromes and discusses the new opportunities and challenges faced with evolving genetic technologies including determining pathogenicity and the utility of large genetic databases. Finally, we will discuss newly described entities that continue the evolving theme of genetic syndromes with phenotypic overlap. Early views that a single genotype associates with a particular phenotype continue to be challenged by our greater understanding of the genotype–phenotype relationship.

### Inherited Arrhythmia Syndromes

#### Long QT Syndrome

Congenital LQTS is diagnosed in the presence of a prolonged corrected QT (QTc) interval after secondary causes (eg, QT-prolonging medications or electrolyte abnormalities) are excluded.<sup>1</sup> The 2013 Heart Rhythm Society/European Heart Rhythm

Association/Asia Pacific Heart Rhythm Association guidelines recommended that LQTS also be diagnosed in the presence of an LQTS risk (Schwartz) score  $\geq 3.5$  or in the presence of an unequivocally pathogenic variant in one of the known LQTS genes.<sup>1</sup> Patients with unexplained syncope and a borderline QTc between 480 and 499 ms in recurrent ECGs, or who were asymptomatic but had repeated measurements of  $>500$  ms, may also be diagnosed with LQTS.<sup>1</sup> The more recent 2015 European Society of Cardiology guidelines softened these criteria to QTc  $\geq 480$  ms in an asymptomatic patient or a QTc  $\geq 460$  ms in the presence of unexplained syncope.<sup>5</sup> It was argued that higher values equated to high-risk LQTS and were, therefore, too conservative.

QT prolongation results from ion channel dysfunction that prolongs cellular repolarization.<sup>6,7</sup> There are currently 15 genes known to cause congenital LQTS; however, the main 3 genotypes account for  $>90\%$  of genetically confirmed LQTS (Table 1).<sup>1,4</sup> The genetic cause remains elusive in 25% of LQTS families.<sup>8</sup> LQT1 is caused by loss-of-function mutations affecting *KCNQ1*, the gene encoding for  $I_{Ks}$  (slow) channel.<sup>4</sup> Both haploinsufficiency and dominant-negative mechanisms of *KCNQ1* mutations are described.<sup>9,10</sup> LQT2 is caused by loss-of-function mutations in *KCNH2*, the gene encoding for the  $I_{Kr}$  (rapid) channel, predominantly because of failure of trafficking to the cell membrane surface.<sup>11,12</sup> Gain-of-function mutations in *SCN5A*, the gene encoding for  $I_{Na^+}$ , causes failed inactivation and increased late current leading to LQT3.<sup>13</sup> Although most LQTS is inherited in an autosomal dominant manner, the rare recessive form (Jervell and Lange-Nielsen syndrome) leads to a severe LQTS phenotype and associated sensorineural deafness.<sup>14</sup> Indeed, greater severity is typical when more than one mutation is identified.<sup>4</sup>

A recent study has suggested that LQTS may be preferentially transmitted from the maternal allele, with higher than expected maternal transmission and inheritance, particularly with LQT1.<sup>15</sup> The authors also found that this phenomenon was not linked to locus-specific grandparental origin allele transmission distortion and in fact seems to be related to the severity of channel dysfunction. The authors postulate that altered potassium channel function may modulate reproduction, particularly *KCNQ1*, which is expressed in the granulosa and trophoblastic cells of the ovaries.<sup>15</sup> LQTS-associated

From the Department of Cardiology, Royal Prince Alfred Hospital, New South Wales, Australia (B.G.); Sydney Medical School, University of Sydney, Australia (B.G.); Agnes Ginges Centre for Molecular Cardiology, Centenary Institute, New South Wales, Australia (B.G.); Cardiology Clinical Academic Group, St George's University of London, United Kingdom (E.R.B.); and St George's University Hospitals NHS Foundation Trust, London, United Kingdom (E.R.B.). Guest Editor for this review was Christopher Semsarian, MBBS, PhD, MPH.

The Data Supplement is available at <http://circgenetics.ahajournals.org/lookup/suppl/doi:10.1161/CIRCGENETICS.116.001571/-/DC1>.

Correspondence to Dr Elijah R. Behr, St George's University of London, Cranmer Terrace, London SW17 0RE, United Kingdom. E-mail ebehr@sgul.ac.uk (*Circ Cardiovasc Genet*. 2016;9:569-577. DOI: 10.1161/CIRCGENETICS.116.001571.)

© 2016 American Heart Association, Inc.

*Circ Cardiovasc Genet* is available at <http://circgenetics.ahajournals.org>

DOI: 10.1161/CIRCGENETICS.116.001571

**Table 1. Genes Associated With Long-QT Syndrome**<sup>84–86,89</sup>

| LQTS Type  | Gene           | Protein       | Current            | Frequency |
|--|----------------|---------------|--------------------|-----------|
| LQT1   | <i>KCNQ1</i>   | Kv7.1         | IKs↓               | 40%–45%   |
| LQT2   | <i>KCNH2</i>   | KV11.1        | IKr↓               | 30%–35%   |
| LQT3   | <i>SCN5A</i>   | Nav1.5        | INa↑               | 10%       |
| LQT4   | <i>ANK2</i>    | Ankyrin-B     | Na+/K+↓            | 1%        |
| LQT5   | <i>KCNE1</i>   | MinK          | IKs↓               | 1%        |
| LQT6   | <i>KCNE2</i>   | MIRP1         | IKr↓               | Rare      |
| LQT7   | <i>KCNJ2</i>   | Kir2.1        | IK1↓               | Rare      |
| LQT8   | <i>CACNA1C</i> | CaV1.2        | ICa-L↑             | Rare      |
| LQT9   | <i>CAV3</i>    | Caveolin 3    | INa↑               | Rare      |
| LQT10  | <i>SCN4B</i>   | SCNβ4 subunit | INa↑               | Rare      |
| LQT11  | <i>AKAP9</i>   | Yotiao        | IKs↓               | Rare      |
| LQT12  | <i>SNTA1</i>   | Syntrophin-α1 | INa↓               | Rare      |
| LQT13  | <i>KCNJ5</i>   | Kir3.4        | IKACH↓             | Rare      |
| LQT14  | <i>CALM1</i>   | Calmodulin 1  | Calcium signalling | Rare      |
| LQT15  | <i>CALM2</i>   | Calmodulin 2  | Calcium signalling | Rare      |
| LQT16  | <i>TRDN</i>    | Triadin       | ICa-L↑             | Rare      |
| Jervell and Lange-Nielsen syndrome (autosomal recessive) |                |               |                    |           |
| JLN1   | <i>KCNQ1</i>   | Kv7.1         | IKs↓               | Rare      |
| JLN2   | <i>KCNE1</i>   | MinK          | IKs↓               | Rare      |

Reprinted from Wilde and Behr.<sup>4</sup>

potassium channel dysfunction has also been proposed recently as a possible mechanism for some stillbirths given that LQTS susceptibility variants can be identified in some cases.<sup>16</sup> This hypothesis follows on from evidence that some sudden infant death syndrome may also be caused by LQTS.<sup>4</sup>

Common variants (single-nucleotide polymorphism or SNPs) at the *NOS1AP* locus have been shown to modify phenotype by affecting both the QT interval and the likelihood of symptoms in LQTS, whereas common variants in *KCNQ1* within the 3′ untranslated region may also influence disease severity.<sup>17–19</sup> A recent genome-wide association and replication study identified a total of 35 SNP-tagged loci associated with QT interval among ≈100 000 individuals of European ancestry, 6 of which were found to have rare coding variants in LQTS patients absent in controls.<sup>20</sup> Although this does not equate to pathogenicity, it is likely that these loci may also modify the LQTS phenotype to some extent.

Genotyping in LQTS is useful diagnostically, therapeutically, and, to a lesser extent, prognostically.<sup>1,8</sup> β-Blockers are the mainstay of therapy, with LQT1 patients, particularly those with mutations in cytoplasmic loops, most protected regardless of drug and LQT2 patients responding best to nadolol.<sup>21,22</sup> Sodium channel blockers (eg, mexiletine) can be useful in addition to β-blockers for patients who have LQT3.<sup>23</sup> High-risk features include severe QT prolongation, boys, female adults, previous cardiac arrest, and syncope despite β-blocker therapy. However, only LQT2 females and those with high-risk genetic profiles (ie, multiple mutations

or Jervell and Lange-Nielsen syndrome) have been associated with sufficient additional risk to suggest a potential risk-stratifying role.<sup>5</sup> Additionally, the biophysical consequences of the mutation may offer alternative methods for stratification.<sup>24</sup>

### Brugada Syndrome

BrS was first formally described in 1992, whereby Brugada et al<sup>25</sup> reported 8 patients with characteristic cove-shaped ST elevation in the right precordial leads with associated sudden cardiac death caused by ventricular fibrillation (VF). Diagnostic criteria from a recent consensus report require the patient to have a spontaneous type 1 Brugada pattern ECG defined as ≥2 mm ST elevation with type 1 morphology in ≥1 right precordial lead V<sub>1</sub> or V<sub>2</sub> in either second, third, or fourth intercostal space.<sup>26</sup> Unlike the earlier 2013 guidelines,<sup>1</sup> if the type 1 pattern is only revealed after sodium channel blocker challenge, the diagnosis of BrS also requires one of the following: documented VF or polymorphic ventricular tachycardia (VT), suspected arrhythmic syncope, family history of sudden cardiac death with negative autopsy, BrS ECG in family members, or nocturnal agonal respiration. This follows recent data suggesting a higher than expected yield of positive results in ajmaline provocation tests in an apparently healthy control population.<sup>27</sup>

Traditionally, BrS has been proposed as a primary electric disease involving a relative impairment of the inward sodium current compared with the transient outward potassium current ( $I_{to}$ ) in the right ventricular outflow tract.<sup>26</sup> However, a transgenic minipig model of a truncating *SCN5A* mutation showed evidence of conduction disease but without a type 1 ECG pattern.<sup>28</sup> This may reflect that haploinsufficiency leads to an isolated conduction disease phenotype without the necessary further genetic variation associated with BrS in humans (see below); the young age of the pigs given that BrS exhibits age-related penetrance; or the absence of  $I_{to}$  in the pig myocardium. The diagnosis of BrS has also required the exclusion of overt structural heart disease and the pathophysiology has been attributed to transmural and epicardial heterogeneity of repolarization. However, there is mounting evidence that BrS may represent one end of a spectrum of subtle structural disease.<sup>29</sup> In particular, imaging studies have suggested abnormalities of the right ventricle and right ventricular outflow tract.<sup>30,31</sup> Epicardial ablation studies have identified fractionated signals consistent with slowed conduction and fibrosis, which have been confirmed by other investigators both invasively and using ECG imaging.<sup>32–34</sup> Subsequent pathological studies in open surgical ablation and sudden death victims from BrS families have confirmed subtle epicardial fibrosis and reduced connexin-43 in the right ventricular outflow tract.<sup>35</sup> An experimental perfused canine wedge model has, however, proposed that these fractionated potentials may result from phase II reentry.<sup>36</sup>

BrS has traditionally been considered as an autosomal dominant inherited disease. However, this convention has also been challenged. The most common associated genetic abnormality is a loss-of-function mutation in *SCN5A*. However, a mutation is identified in only one fifth of patients.<sup>4</sup> Strong linkage studies associating *SCN5A* with BrS do not exist, with

the exception of families with overlap phenotypes with LQTS and PCCD.<sup>37</sup> Additionally, the genotype–phenotype correlation within families shows worrying mismatch. Probst et al<sup>38</sup> described 5 out of 13 large BrS families with known *SCN5A* mutations with several individuals with a type 1 Brugada ECG pattern within the same family who were mutation negative. Mutations in 22 other genes-encoding components and interacting proteins of the sodium channel, calcium channel, and potassium channels have also been implicated in BrS (Table 2); however, few are based on strong linkage studies, and most are rare.<sup>4,39</sup>

Furthermore, a genome-wide association study identified 3 SNPs with additive effects on the likelihood of BrS irrespective of the presence of an *SCN5A* mutation.<sup>40</sup> These included 2 SNPs at the *SCN5A* and *SCN10A* locus; *SCN10A* encoding the  $\alpha$  subunit of the neuronal Nav1.8 current associated with nociception. This prompted investigation by Hu et al<sup>41</sup> who suggested that mutations in *SCN10A* are present in up to 16% of BrS patients. A contemporaneous study assessed the burden of rare coding variants in BrS-associated genes in patients compared with a control group and only showed enrichment of *SCN5A*. This concluded that previously reported genes including *SCN10A* do not account for BrS cases in patients of European ancestry.<sup>42</sup> Additionally, a subsequent study of 45 known cardiac genes in *SCN5A*-negative cases showed enrichment of the *DSC2* gene linked to arrhythmogenic right ventricular cardiomyopathy but not the other BrS genes.<sup>43</sup> These data reinforced the possible role for arrhythmogenic right ventricular cardiomyopathy-associated rare variants in BrS that had already been proposed for the *PKP2* gene in an earlier study.<sup>44</sup> Another study of potential novel candidate genes in *SCN5A*-negative cases showed a similar low yield although a probable mutation was identified in *TBX5*, a transcription factor for *SCN5A* and *SCN10A*.<sup>45</sup> The study also confirmed previous findings that SNPs at the *SCN10A* locus associates with BrS, but rare variation was not significantly associated. Collectively, these studies support the theory that BrS is an oligogenic disease with the potential that ion channel and structural rare variants may have modulatory influences on a common substrate. Together with other factors such as age, sex, and environmental exposures, these affect the susceptibility for the BrS phenotype and potentially contribute to the subtle structural disease that has been associated with the disorder. This may also account for the unexpectedly high yield of ajmaline provocation in the general population, whereby susceptibility to BrS is partly because of genetic variation prevalent in the general population.<sup>27</sup> The role for diagnostic genetic testing is, therefore, relatively limited at this time until we are able to dissect the interaction of rare and common variation.

### Catecholaminergic Polymorphic Ventricular Tachycardia

CPVT is a highly lethal, rare inherited arrhythmia syndrome characterized by bidirectional and polymorphic VT, which is adrenergically stimulated, typically through physical exertion.<sup>46,47</sup> The diagnostic criteria require a structurally normal heart and normal baseline ECG, with the development of polymorphic or bidirectional VT after exercise or catecholamine,

usually in a patient aged <40 years.<sup>1</sup> CPVT can also be diagnosed in patients who are found to be carriers of a known clearly pathogenic genetic variant and in family members of an index case who develop premature ventricular contractions during exercise.<sup>1</sup>

The pathophysiology of CPVT is dependent on abnormal intracellular calcium handling leading to triggered activity.<sup>48</sup> Linkage studies in 2 large Finnish families with autosomal dominantly inherited CPVT first identified linkage at chromosome 1q42-43.<sup>49</sup> Subsequent studies indicated the gene encoding the cardiac ryanodine receptor gene (*RYR2*) to be responsible (CPVT1).<sup>1,4</sup> In CPVT, *RYR2* is leaky under adrenergic activation, causing an excess of calcium release into the cytosol in diastole, resulting in increased activation of the sodium–calcium exchanger and delayed after-depolarizations.<sup>50</sup> A rarer form of autosomal recessive CPVT (CPVT2) is known to result from mutations in the cardiac calsequestrin gene (*CASQ2*) an important calcium storage and buffering protein in the sarcoplasmic reticulum interacting with *RYR2*.<sup>4</sup> A

**Table 2. Genes Associated With Brugada Syndrome**

| Name  | Gene            | Protein   | Prevalence |
|-------|-----------------|---|------------|
| BrS1  | <i>SCN5A</i>    | $\alpha$ -subunit Nav1.5 Sodium channel                       | 20%–25%    |
| BrS2  | <i>GPD1L</i>    | Glycerol-3-phosphate dehydrogenase 1-like                     | Rare       |
| BrS3  | <i>CACNA1C</i>  | $\alpha$ -subunit $\alpha$ 1C Cav1.2 Calcium channel          | 1%–2%      |
| BrS4  | <i>CACNB2b</i>  | $\beta$ -subunit Cav $\beta$ 2b calcium channel               | 1%–2%      |
| BrS5  | <i>SCN1b</i>    | $\beta$ -subunit Nav $\beta$ 1 sodium channel                 | Rare       |
| BrS6  | <i>KCNE3</i>    | $\beta$ -subunit MiRP2 potassium channel                      | Rare       |
| BrS7  | <i>SCN3b</i>    | $\beta$ -subunit Nav $\beta$ 3 sodium channel                 | Rare       |
| BrS8  | <i>HCN4</i>     | Hyperpolarization-activated cyclic nucleotide-gated channel 4 | Rare       |
| BrS9  | <i>KCND3</i>    | $\alpha$ -subunit KV4.3 potassium channel                     | Rare       |
| BrS10 | <i>KCNJ8</i>    | $\alpha$ -subunit KIR6.1 potassium channel                    | Rare       |
| BrS11 | <i>CACNA2D1</i> | $\delta$ -subunit Cav $\alpha$ 2 $\delta$ 1 calcium channel   | Rare       |
| BrS12 | <i>KCNE5</i>    | $\beta$ -subunit potassium channel                            | Rare       |
| BrS13 | <i>RANGRF</i>   | RAN guanine nucleotide release factor                         | Rare       |
| BrS14 | <i>KCND2</i>    | $\alpha$ -subunit KV4.2 potassium channel                     | Rare       |
| BrS15 | <i>TRPM4</i>    | Calcium-activated nonselective ion channel                    | Rare       |
| BrS16 | <i>SCN2B</i>    | $\beta$ -subunit Nav $\beta$ 2 sodium channel                 | Rare       |
| BrS17 | <i>PKP2</i>     | Plakophilin 2   | Rare       |
| BrS18 | <i>ABCC9</i>    | ATP-sensitive potassium channels                              | Rare       |
| BrS19 | <i>SLMAP</i>    | Sarcolemma-associated protein                                 | Rare       |
| BrS20 | <i>KCNH2</i>    | $\alpha$ -subunit of HERG potassium channel                   | Rare       |
| BrS21 | <i>SCN10A</i>   | $\alpha$ -subunit Nav1.8 sodium channel                       | 1%–16%     |
| BrS22 | <i>FGF12</i>    | Fibroblast growth factor 12                                   | Rare       |
| BrS23 | <i>SEMA3A</i>   | Semaphorin family protein                                     | Rare       |

Reprinted from Wilde and Behr<sup>4</sup> and Gourraud et al.<sup>39</sup>

recent study has also shown linkage to a heterozygous *CASQ2* variant in autosomal dominant CPVT for the first time.<sup>51</sup>

Collectively, these 2 genes combined account for ≈50% to 65% of cases of CPVT.<sup>8</sup> Other genes, including calmodulin (*CALM1*), triadin (*TRDN*), *TECLR*, ankyrin-B (*ANK2*), and *KCNJ2* account for rare cases of CPVT.<sup>4,52</sup> Whether *ANK2* and *KCNJ2* represent phenocopies or overlap syndromes with LQT4 and 7, respectively, remains a moot point (Table 3).

### Short-QT Syndrome

SQTS is a rare inherited arrhythmia syndrome characterized by syncope or sudden cardiac death associated with a shortened QTc interval. The diagnosis requires the presence of a very short QTc ≤330 ms or a QTc ≤360 ms in association with one of the following: family history of SQTS, pathogenic mutation, family history of sudden death at age ≤40 years, or survival of a VT/VF episode in the absence of heart disease.<sup>1</sup> SQTS was first described in 2000,<sup>53</sup> and since then, candidate gene studies have linked SQTS to missense mutations causing gain of function in potassium channels (*KCNH2*-SQT1, *KCNQ1*-SQT2, and *KCNJ2*-SQT3).<sup>4,54</sup> There is also an association with loss-of-function mutations in calcium channels (*CACNA1C* and *CACNB2b*) in association with a BrS phenotype.<sup>4,54</sup> SQTS is rare with the largest series of 73 patients demonstrating a low genetic yield of 14% despite familial disease being evident in 44%.<sup>54</sup> It is unclear whether there may be an overlap with IVF cases with only moderate QT shortening leading to underdiagnosis.

### Idiopathic Ventricular Fibrillation

IVF describes patients who experience a resuscitated cardiac arrest, ideally with documented VF in the absence of other causes for VF, that is, metabolic, toxicological, cardiac (including other channelopathies and structural heart disease), respiratory, and infectious.<sup>1</sup> The arrhythmias are believed to be precipitated by short-coupled premature ventricular contractions due to Purkinje fiber potentials leading to polymorphic VT.<sup>55–57</sup>

Although IVF remains largely unexplained, the genetic basis for disease has been proposed in some cases. Alders et al<sup>58</sup> linked disease with *DPP6* gene though a genome-wide haplotype-sharing analysis in 3 distantly related Dutch

families. Carriers demonstrated high penetrance with 50% experiencing an episode of VF or sudden death before age 58 years. Marsman et al<sup>59</sup> used exome sequencing to link IVF to *CALM1* in a family of Moroccan descent. A recent report described a family with a likely IVF phenotype, with short-coupled premature ventricular contractions at rest and a novel *RYR2* mutation H29D causing diastolic calcium leak at rest.<sup>60</sup> Finally, a recent report highlights *IRX3* as a putative gene in humans and animal models.<sup>61</sup> Collectively, these reports suggest potential monogenic causes for IVF. However, due to the lack of large IVF families, it remains possible that the disease may be oligogenic.<sup>62</sup> Additionally, the prevalence of IVF is declining as more patients initially diagnosed with IVF are subsequently diagnosed with a different inherited arrhythmia syndrome such as BrS.<sup>62</sup>

### Progressive Cardiac Conduction System Disease

PCCD is diagnosed in young patients (<50 years of age) exhibiting abnormalities of conduction with and without a structurally normal heart in the absence of skeletal myopathies.<sup>1</sup> When there is a clear familial pattern to disease, it is typically autosomal dominant, and genetic testing has diagnostic utility.<sup>1</sup> The majority of familial cases with a structurally normal heart are attributed to mutations in *SCN5A* and *TRPM4*.<sup>8</sup> Cases because of *SCN5A* mutations can be associated with a BrS overlap syndrome.<sup>8,37,63</sup> There are also rare associations with other genes (*SCN1B*, *SCN10A*, and *KCNK17*).<sup>64</sup> PCCD with congenital heart disease is associated with transcription factor mutations (eg, *NKX2.5* and *GATA4*)<sup>65</sup> and PCCD with dilated cardiomyopathy or LV dysfunction is associated with lamin A/C (*LMNA*) or desmin (*DES*) variants.<sup>8</sup> The presence of a *LMNA* mutation carries prognostic utility and may influence ICD implantation.<sup>1</sup>

### Genetic Testing and Next-Generation Sequencing

Identification of the likely pathogenic variant in a family permits predictive (cascade) testing in family members. Those who are not found to carry the variant are released from regular clinical screening.<sup>66</sup> The best approach to improving the pretest probability of a pathogenic variant being identified is to ensure that the clinical phenotype is clearly defined, whether there is a family history of disease and that only genes that are plausible for the phenotype are assessed.<sup>8</sup>

Diagnostic testing had traditionally relied on Sanger sequencing, and disease causation or pathogenicity was often inferred only by absence of a variant in relatively small control populations. Since 2005, next-generation sequencing methods have become increasingly available including cardiac gene panel testing, whole-exome sequencing, and whole-genome sequencing. These allow large volumes of data to be analyzed quickly and comprehensively at lower cost and have helped identify causative genes in disease where traditional methods have previously failed.<sup>67</sup> These differing approaches and their relevant advantages and disadvantages are shown in the [Data Supplement](#).

Next-generation sequencing has also facilitated large public databases of genetic variation such as the 1000 Genomes project, the Exome Sequencing Project, or Exome Aggregation Consortium that have indicated rare variants to be more

**Table 3. Genes Associated With CPVT**

| Name           | Gene         | Protein  | Frequency |
|----------------|--------------|--|-----------|
| CPVT1          | <i>RYR2</i>  | Cardiac ryanodine receptor 2   | 50%–60%   |
| CPVT2          | <i>CASQ2</i> | Cardiac calsequestrin  | ≈5%       |
| CPVT3          | <i>TECLR</i> | Originally mapped to chromosome 7 p14-p22, now reallocated to chromosome 4 | Rare      |
| CPVT4          | <i>CALM1</i> | Calmodulin   | Rare      |
| CPVT5          | <i>TRDN</i>  | Triadin  | Rare      |
| ? LQT4 overlap | <i>ANK2</i>  | Ankyrin B  | Rare      |
| ? LQT7 overlap | <i>KCNJ2</i> | Potassium inwardly rectifying channel Kir2.1                               | Rare      |

Reprinted from Wilde and Behr.<sup>4</sup>

common than expected.<sup>26</sup> For example, in *SCN5A*, there is a 3% to 5% background noise of rare variants among healthy individuals.<sup>68</sup> Therefore,  $\leq 1$  in 20 so-called positive genetic tests for BrS could in fact be false positives.<sup>26</sup> The current signal-to-noise ratio for the top 4 inherited arrhythmia genes ranges between 4.5:1 (*SCN5A* for LQT3) and 70:1 (*KCNQ1* for LQT1).<sup>69</sup> Additionally, variants previously thought of as being pathogenic in early Sanger studies are now accepted to be too common to be causative of rare disease and are, therefore, false positives. For example, 1 in 23 individuals in the Exome Sequencing Project database carried a previously published BrS-associated variant.<sup>70</sup> Similarly Refsgaard et al<sup>71</sup> identified 33 rare variants previously associated with LQTS in the Exome Sequencing Project database, suggesting a disease prevalence of 1 in 31 rather than 1 in 2000. Ghouse et al<sup>72</sup> showed that 10 variants previously associated with LQTS were in fact found frequently among a Danish population sample. Finally, Van Driest et al<sup>73</sup> showed no significant difference in arrhythmic phenotype between those with and without rare *KCNH2* or *SCN5A* variants in an unselected population. They also showed poor concordance among laboratories when designating rare variants as pathogenic.

Thus, to strengthen utility of contemporary genetic testing, there are stringent criteria to determine pathogenicity. It is no longer considered a binary variable but instead a probabilistic spectrum of pathogenicity as shown in Figure 1.<sup>66,74,75</sup> Criteria have been published by the American College of Medical Genetics and Genomics and are currently being updated with cardiac-specific recommendations.<sup>76</sup>

### Assessing Pathogenicity

Online databases of previously reported pathogenic variants have provided important tools for recognizing unequivocally pathogenic variants although they are only as robust as the data entered and require regular updating and stringent curation. The ClinVar ([www.ncbi.nlm.nih.gov/clinvar/](http://www.ncbi.nlm.nih.gov/clinvar/)) and ClinGen ([www.clinicalgenome.org](http://www.clinicalgenome.org)) initiatives offer important opportunities for global harmonization of such data.<sup>77</sup>

Rarity can still be helpful when assessing likelihood of pathogenicity.<sup>78</sup> The public databases are integral to determining which variants are truly rare or novel and which are part of normal background genetic variation within cardiac inherited arrhythmia genes.<sup>26</sup> The prevalence of the disease being tested for should, however, be considered. For example, the disease


prevalence of CPVT is 1/10000; therefore, a variant with a population frequency of 0.0001, while rare, would potentially account for all CPVT cases globally and is, therefore, too common to be pathogenic. Novelty or substantially lower allele frequency than the disease prevalence is, therefore, required to support pathogenicity.

Loss of function is a known mechanism of disease pathogenesis for most inherited arrhythmia syndromes and an important step in assessing pathogenicity is reviewing the predicted effect of the variant on protein function. This is relatively straightforward for nonsense, frameshift or splice-site mutations but less so for missense variants. In silico software tools and functional studies, either in vivo or in vitro, can then also be used to assess the physiological effect of the variant and predict likelihood of pathogenicity although they are not always reliable.<sup>26,76,78</sup> Kapplinger et al<sup>68</sup> used multiple in silico tools and ion channel topography to improve variant classification as benign or pathogenic. All tools (bar the Grantham score) showed some predictive value; although none showed strong independent utility.

Despite these efforts, variants of uncertain significance have become increasingly problematic. These are typically absent from population databases and may have computational evidence suggestive of pathogenicity. Lack of functional or segregation data, however, means that the variant cannot be classified as likely pathogenic or benign. Genetic counseling is critical to ensure the patients understand this potential complication of genetic testing.<sup>8</sup> Predictive testing should, therefore, only be performed in monogenic disease when a variant has been classified as pathogenic or likely pathogenic.<sup>66</sup>

### Changing Landscape of Inherited Arrhythmia Syndromes

Inherited arrhythmia syndromes have typically been attributed to autosomal dominant inheritance because of monogenic abnormalities. As discussed above, these diseases may be genetically more complex and oligogenic, particularly in BrS (Figure 2A). The simple genotype equals phenotype equation is also increasingly questioned. This was first evidenced by *SCN5A* overlap syndrome of BrS, PCCD, and LQTS<sup>37</sup> followed by association with other phenotypes,<sup>63</sup> as well as dilated cardiomyopathy and atrial fibrillation.<sup>4</sup> There is also evidence of CPVT and LQTS overlap phenotypes because of different *ANK2* and *KCNJ2* mutations. The



**Figure 1.** Probabilistic nature of genetic results. VUS indicates variants of uncertain significance. Reprinted from Maron et al<sup>74</sup> with permission of the publisher.



in a cohort of 33 additional unaffected patients with gene-negative LQTS revealed 4 additional patients with homozygous or compound heterozygous *TRDN* frameshift mutations. All patients carried the same distinct phenotype of extensive T-wave inversion in the precordial ECG leads V<sub>1</sub> through V<sub>4</sub> (similar to arrhythmogenic right ventricular cardiomyopathy), QTc prolongation, and severe disease expression at a young age, with arrhythmias particularly precipitated by exercise.<sup>89</sup> The mechanism underpinning the *TRDN* knockout syndrome seems to be reduction in negative feedback on the L-type calcium channel leading to increased intracellular calcium, increased frequency of sarcoplasmic reticulum calcium release and subsequent VT, particularly with adrenergic stimulation.<sup>88</sup> L-type calcium channel-blocking agents have been proposed as potential therapeutic options in patients with this distinct syndrome.<sup>89</sup> The triadin knockout syndrome seems to be an overlap syndrome with a LQTS/CPVT/arrhythmogenic right ventricular cardiomyopathy overlap phenotype (Figure 2B).

### Gene Therapies for Inherited Arrhythmia Syndromes

Recent work has highlighted that it may be possible to cure arrhythmia syndromes in the not too distant future. A *CASQ2* knockout mouse model of CPVT was rescued with *CASQ2* via an adeno-associated viral vector (AAV9). The arrhythmic phenotype and ultrastructural abnormalities were also rescued with absence of arrhythmias in the rescued mice.<sup>90</sup> Subsequent work showed similar success in rescuing the phenotype from a knock-in murine model with *CASQ2* R33Q homozygous mutation with administration of AAV9-*CASQ2* at birth. The mice continued to display no evidence of CPVT at 1 year of age. Although still early in evolution, these animal models provide an exciting insight into where potential gene therapy options may arise.

### Conclusions

Our understanding of inherited arrhythmia syndromes continues to evolve. Clinicians must continue to use caution when interpreting genetic variants and when using genetic results to determine management of patients and families with inherited arrhythmia syndromes. There is increasing genetic and phenotypic heterogeneity and mounting evidence that BrS is an oligogenic disorder. The era of next-generation sequencing has evolved rapidly providing challenges including attributing pathogenicity, background genetic noise, and increased detection of variants of uncertain significance; as well as new opportunities, including identification of new inherited arrhythmia entities such as the calmodulinopathies and Triadin disease. Our early understanding that a particular genotype causes a specific phenotype is further challenged with these new entities.

### Disclosures

None.

### References

- Priori SG, Wilde AA, Horie M, Cho Y, Behr ER, Berul C, et al. HRS/EHRA/APHS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes: document endorsed by HRS, EHRA, and APHS in May 2013 and by ACCF, AHA, PACES, and AEPC in June 2013. *Heart Rhythm*. 2013;10:1932–1963. doi: 10.1016/j.hrthm.2013.05.014.
- Behr ER, Dalageorgou C, Christiansen M, Syrris P, Hughes S, Tome Esteban MT, et al. Sudden arrhythmic death syndrome: familial evaluation identifies inheritable heart disease in the majority of families. *Eur Heart J*. 2008;29:1670–1680. doi: 10.1093/eurheartj/ehn219.
- Miles CJ, Behr ER. The role of genetic testing in unexplained sudden death. *Transl Res*. 2016;168:59–73. doi: 10.1016/j.trsl.2015.06.007.
- Wilde AA, Behr ER. Genetic testing for inherited cardiac disease. *Nat Rev Cardiol*. 2013;10:571–583. doi: 10.1038/nrcardio.2013.108.
- Priori SG, Blomström-Lundqvist C, Mazzanti A, Blom N, Borggrefe M, Camm J, et al. 2015 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: the Task Force for the Management of Patients with Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death of the European Society of Cardiology (ESC). Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC). *Eur Heart J*. 2015;36:2793–2867. doi: 10.1093/eurheartj/ehv316.
- Keating MT, Sanguinetti MC. Molecular genetic insights into cardiovascular disease. *Science*. 1996;272:681–685.
- Chiang CE, Roden DM. The long QT syndromes: genetic basis and clinical implications. *J Am Coll Cardiol*. 2000;36:1–12.
- Ackerman MJ, Priori SG, Willems S, Berul C, Brugada R, Calkins H, et al. HRS/EHRA expert consensus statement on the state of genetic testing for the channelopathies and cardiomyopathies this document was developed as a partnership between the Heart Rhythm Society (HRS) and the European Heart Rhythm Association (EHRA). *Heart Rhythm*. 2011;8:1308–1339. doi: 10.1016/j.hrthm.2011.05.020.
- Heijman J, Spätjens RL, Seyen SR, Lentink V, Kuijpers HJ, Boulet IR, et al. Dominant-negative control of cAMP-dependent IKs upregulation in human long-QT syndrome type 1. *Circ Res*. 2012;110:211–219. doi: 10.1161/CIRCRESAHA.111.249482.
- Schwartz PJ, Ackerman MJ, George AL Jr, Wilde AA. Impact of genetics on the clinical management of channelopathies. *J Am Coll Cardiol*. 2013;62:169–180. doi: 10.1016/j.jacc.2013.04.044.
- Sanguinetti MC, Jiang C, Curran ME, Keating MT. A mechanistic link between an inherited and an acquired cardiac arrhythmia: HERG encodes the IKr potassium channel. *Cell*. 1995;81:299–307.
- Curran ME, Splawski I, Timothy KW, Vincent GM, Green ED, Keating MT. A molecular basis for cardiac arrhythmia: HERG mutations cause long QT syndrome. *Cell*. 1995;80:795–803.
- Wang Q, Shen J, Splawski I, Atkinson D, Li Z, Robinson JL, et al. SCN5A mutations associated with an inherited cardiac arrhythmia, long QT syndrome. *Cell*. 1995;80:805–811.
- Jervell A, Lange-Nielsen F. Congenital deaf-mutism, functional heart disease with prolongation of the Q-T interval and sudden death. *Am Heart J*. 1957;54:59–68.
- Itoh H, Berthet M, Fressart V, Denjoy I, Maugendre S, Klug D, et al. Asymmetry of parental origin in long QT syndrome: preferential maternal transmission of KCNQ1 variants linked to channel dysfunction. *Eur J Hum Genet*. 2016;24:1160–1166. doi: 10.1038/ejhg.2015.257.
- Crotti L, Tester DJ, White WM, Bartos DC, Insolia R, Besana A, et al. Long QT syndrome-associated mutations in intrauterine fetal death. *JAMA*. 2013;309:1473–1482. doi: 10.1001/jama.2013.3219.
- Crotti L, Monti MC, Insolia R, Peljto A, Goosen A, Brink PA, et al. NOS1AP is a genetic modifier of the long-QT syndrome. *Circulation*. 2009;120:1657–1663. doi: 10.1161/CIRCULATIONAHA.109.879643.
- Tomás M, Napolitano C, De Giuli L, Bloise R, Subirana I, Malovini A, et al. Polymorphisms in the NOS1AP gene modulate QT interval duration and risk of arrhythmias in the long QT syndrome. *J Am Coll Cardiol*. 2010;55:2745–2752. doi: 10.1016/j.jacc.2009.12.065.
- Amin AS, Giudicessi JR, Tijssen AJ, Spanjaart AM, Reckman YJ, Kl-emens CA, et al. Variants in the 3' untranslated region of the KCNQ1-encoded Kv7.1 potassium channel modify disease severity in patients with type 1 long QT syndrome in an allele-specific manner. *Eur Heart J*. 2012;33:714–723. doi: 10.1093/eurheartj/ehr473.
- Arking DE, Pulit SL, Crotti L, van der Harst P, Munroe PB, Koopmann TT, et al; CARE Consortium; COGENT Consortium; DCCT/EDIC; eMERGE Consortium; HRGEN Consortium. Genetic association study of QT interval highlights role for calcium signaling pathways in myocardial repolarization. *Nat Genet*. 2014;46:826–836. doi: 10.1038/ng.3014.
- Abu-Zeitone A, Peterson DR, Polonsky B, McNitt S, Moss AJ. Efficacy of different beta-blockers in the treatment of long QT syndrome. *J Am Coll Cardiol*. 2014;64:1352–1358. doi: 10.1016/j.jacc.2014.05.068.

22. Barsheshet A, Goldenberg I, O-Uchi J, Moss AJ, Jons C, Shimizu W, et al. Mutations in cytoplasmic loops of the KCNQ1 channel and the risk of life-threatening events: implications for mutation-specific response to  $\beta$ -blocker therapy in type 1 long-QT syndrome. *Circulation*. 2012;125:1988–1996. doi: 10.1161/CIRCULATIONAHA.111.048041.
23. Mazzanti A, Maragna R, Faragli A, Monteforte N, Bloise R, Memmi M, et al. Gene-specific therapy with mexiletine reduces arrhythmic events in patients with long QT syndrome type 3. *J Am Coll Cardiol*. 2016;67:1053–1058. doi: 10.1016/j.jacc.2015.12.033.
24. Jons C, O-Uchi J, Moss AJ, Reumann M, Rice JJ, Goldenberg I, et al. Use of mutant-specific ion channel characteristics for risk stratification of long QT syndrome patients. *Sci Transl Med*. 2011;3:76ra28.
25. Brugada P, Brugada J. Right bundle branch block, persistent ST segment elevation and sudden cardiac death: a distinct clinical and electrocardiographic syndrome. A multicenter report. *J Am Coll Cardiol*. 1992;20:1391–1396.
26. Antzelevitch C, Yan GX, Ackerman MJ, Borggreffe M, Corrado D, Guo J, et al. J-Wave syndromes expert consensus conference report: emerging concepts and gaps in knowledge. *Heart Rhythm*. 2016;13:e295–324.
27. Hasdemir C, Payzin S, Kocabas U, Sahin H, Yildirim N, Alp A, et al. High prevalence of concealed Brugada syndrome in patients with atrioventricular nodal reentrant tachycardia. *Heart Rhythm*. 2015;12:1584–1594. doi: 10.1016/j.hrthm.2015.03.015.
28. Park DS, Cerrone M, Morley G, Vasquez C, Fowler S, Liu N, et al. Genetically engineered SCN5A mutant pig hearts exhibit conduction defects and arrhythmias. *J Clin Invest*. 2015;125:403–412. doi: 10.1172/JCI76919.
29. Gray B, Semsarian C, Sy RW. Brugada syndrome: a heterogeneous disease with a common ECG phenotype? *J Cardiovasc Electrophysiol*. 2014;25:450–456. doi: 10.1111/jce.12366.
30. Catalano O, Antonaci S, Moro G, Mussida M, Frascaroli M, Baldi M, et al. Magnetic resonance investigations in Brugada syndrome reveal unexpectedly high rate of structural abnormalities. *Eur Heart J*. 2009;30:2241–2248. doi: 10.1093/eurheartj/ehp252.
31. Bastiaenen R, Cox A, Castelletti S, Wijeyeratne Y, Colbeck N, Pakroo N, et al. Late gadolinium enhancement in Brugada syndrome: a marker for subtle underlying cardiomyopathy? *Heart Rhythm*. In press.
32. Nademanee K, Veerakul G, Chandanamattha P, Chaothawee L, Ariyachai-panich A, Jirasirojanakorn K, et al. Prevention of ventricular fibrillation episodes in Brugada syndrome by catheter ablation over the anterior right ventricular outflow tract epicardium. *Circulation*. 2011;123:1270–1279. doi: 10.1161/CIRCULATIONAHA.110.972612.
33. Brugada J, Pappone C, Berruezo A, Vicedomini G, Manguso F, Ciconte G, et al. Brugada syndrome phenotype elimination by epicardial substrate ablation. *Circ Arrhythm Electrophysiol*. 2015;8:1373–1381. doi: 10.1161/CIRCEP.115.003220.
34. Zhang J, Sacher F, Hoffmayer K, O'Hara T, Strom M, Cuculich P, et al. Cardiac electrophysiological substrate underlying the ECG phenotype and electrogram abnormalities in Brugada syndrome patients. *Circulation*. 2015;131:1950–1959. doi: 10.1161/CIRCULATIONAHA.114.013698.
35. Nademanee K, Raju H, de Noronha SV, Papadakis M, Robinson L, Rothery S, et al. Fibrosis, connexin-43, and conduction abnormalities in the Brugada syndrome. *J Am Coll Cardiol*. 2015;66:1976–1986. doi: 10.1016/j.jacc.2015.08.862.
36. Szél T, Antzelevitch C. Abnormal repolarization as the basis for late potentials and fractionated electrograms recorded from epicardium in experimental models of Brugada syndrome. *J Am Coll Cardiol*. 2014;63:2037–2045. doi: 10.1016/j.jacc.2014.01.067.
37. Bezzina C, Veldkamp MW, van Den Berg MP, Postma AV, Rook MB, Viersma JW, et al. A single Na<sup>+</sup> channel mutation causing both long-QT and Brugada syndromes. *Circ Res*. 1999;85:1206–1213.
38. Probst V, Wilde AA, Barc J, Sacher F, Babuty D, Mabo P, et al. SCN5A mutations and the role of genetic background in the pathophysiology of Brugada syndrome. *Circ Cardiovasc Genet*. 2009;2:552–557. doi: 10.1161/CIRCGENETICS.109.853374.
39. Gourraud JB, Barc J, Thollet A, Le Scouarnec S, Le Marec H, Schott JJ, et al. The Brugada syndrome: a rare arrhythmia disorder with complex inheritance. *Front Cardiovasc Med*. 2016;3:9. doi: 10.3389/fcvm.2016.00009.
40. Bezzina CR, Barc J, Mizusawa Y, Remme CA, Gourraud JB, Simonet F, et al. Common variants at SCN5A-SCN10A and HEY2 are associated with Brugada syndrome, a rare disease with high risk of sudden cardiac death. *Nat Genet*. 2013;45:1044–1049. doi: 10.1038/ng.2712.
41. Hu D, Barajas-Martinez H, Pfeiffer R, Dezi F, Pfeiffer J, Buch T, et al. Mutations in SCN10A are responsible for a large fraction of cases of Brugada syndrome. *J Am Coll Cardiol*. 2014;64:66–79. doi: 10.1016/j.jacc.2014.04.032.
42. Le Scouarnec S, Karakachoff M, Gourraud JB, Lindenbaum P, Bonnaud S, Portero V, et al. Testing the burden of rare variation in arrhythmia-susceptibility genes provides new insights into molecular diagnosis for Brugada syndrome. *Hum Mol Genet*. 2015;24:2757–2763. doi: 10.1093/hmg/ddv036.
43. Di Resta C, Pietrelli A, Sala S, Della Bella P, De Bellis G, Ferrari M, et al. High-throughput genetic characterization of a cohort of Brugada syndrome patients. *Hum Mol Genet*. 2015;24:5828–5835. doi: 10.1093/hmg/ddv302.
44. Cerrone M, Lin X, Zhang M, Agullo-Pascual E, Pfenniger A, Chkourko Gusky H, et al. Missense mutations in plakophilin-2 cause sodium current deficit and associate with a Brugada syndrome phenotype. *Circulation*. 2014;129:1092–1103. doi: 10.1161/CIRCULATIONAHA.113.003077.
45. Behr ER, Savio-Galimberti E, Barc J, Holst AG, Petropoulou E, Prins BP, et al; UK10K Consortium. Role of common and rare variants in SCN10A: results from the Brugada syndrome QRS locus gene discovery collaborative study. *Cardiovasc Res*. 2015;106:520–529. doi: 10.1093/cvr/cvv042.
46. Leenhardt A, Lucet V, Denjoy I, Grau F, Ngoc DD, Coumel P. Catecholaminergic polymorphic ventricular tachycardia in children. A 7-year follow-up of 21 patients. *Circulation*. 1995;91:1512–1519.
47. Coumel P, Fidelle J, Lucet V, Attuel P, Bouvrain Y. Catecholamine-induced severe ventricular arrhythmias with Adams-Stokes syndrome in children: report of four cases. *Br Heart J*. 1978;40:28–37.
48. Priori SG, Chen SR. Inherited dysfunction of sarcoplasmic reticulum Ca<sup>2+</sup> handling and arrhythmogenesis. *Circ Res*. 2011;108:871–883. doi: 10.1161/CIRCRESAHA.110.226845.
49. Swan H, Piippo K, Viitasalo M, Heikkilä P, Paavonen T, Kainulainen K, et al. Arrhythmic disorder mapped to chromosome 1q42-q43 causes malignant polymorphic ventricular tachycardia in structurally normal hearts. *J Am Coll Cardiol*. 1999;34:2035–2042.
50. Jiang D, Wang R, Xiao B, Kong H, Hunt DJ, Choi P, et al. Enhanced store overload-induced Ca<sup>2+</sup> release and channel sensitivity to luminal Ca<sup>2+</sup> activation are common defects of RyR2 mutations linked to ventricular tachycardia and sudden death. *Circ Res*. 2005;97:1173–1181. doi: 10.1161/01.RES.0000192146.85173.4b.
51. Gray B, Bagnall RD, Lam L, Ingles J, Turner C, Haan E, et al. A novel heterozygous mutation in cardiac calsequestrin causes autosomal dominant catecholaminergic polymorphic ventricular tachycardia. *Heart Rhythm*. 2016;13:1652–1660. doi: 10.1016/j.hrthm.2016.05.004.
52. Lieve KV, van der Werf C, Wilde AA. Catecholaminergic polymorphic ventricular tachycardia. *Circ J*. 2016;80:1285–1291. doi: 10.1253/circj.CJ-16-0326.
53. Gussak I, Brugada P, Brugada J, Wright RS, Kopecky SL, Chaitman BR, et al. Idiopathic short QT interval: a new clinical syndrome? *Cardiology*. 2000;94:99–102. doi: 47299.
54. Mazzanti A, Kanthan A, Monteforte N, Memmi M, Bloise R, Novelli V, et al. Novel insight into the natural history of short QT syndrome. *J Am Coll Cardiol*. 2014;63:1300–1308. doi: 10.1016/j.jacc.2013.09.078.
55. Haïssaguerre M, Shoda M, Jaïs P, Nogami A, Shah DC, Kautzner J, et al. Mapping and ablation of idiopathic ventricular fibrillation. *Circulation*. 2002;106:962–967.
56. Viskin S, Lesh MD, Eldar M, Fish R, Setbon I, Laniado S, et al. Mode of onset of malignant ventricular arrhythmias in idiopathic ventricular fibrillation. *J Cardiovasc Electrophysiol*. 1997;8:1115–1120.
57. Aizawa Y, Chinushi M, Hasegawa K, Naiki N, Horie M, Kaneko Y, et al. Electrical storm in idiopathic ventricular fibrillation is associated with early repolarization. *J Am Coll Cardiol*. 2013;62:1015–1019. doi: 10.1016/j.jacc.2013.05.030.
58. Alders M, Koopmann TT, Christiaans I, Postema PG, Beekman L, Tanck MW, et al. Haplotype-sharing analysis implicates chromosome 7q36 harboring DPP6 in familial idiopathic ventricular fibrillation. *Am J Hum Genet*. 2009;84:468–476. doi: 10.1016/j.ajhg.2009.02.009.
59. Marsman RF, Barc J, Beekman L, Alders M, Dooijes D, van den Wijngaard A, et al. A mutation in CALM1 encoding calmodulin in familial idiopathic ventricular fibrillation in childhood and adolescence. *J Am Coll Cardiol*. 2014;63:259–266. doi: 10.1016/j.jacc.2013.07.091.
60. Cheung JW, Meli AC, Xie W, Mittal S, Reiken S, Wronska A, et al. Short-coupled polymorphic ventricular tachycardia at rest linked to a novel ryanodine receptor (RyR2) mutation: leaky RyR2 channels under non-stress conditions. *Int J Cardiol*. 2015;180:228–236. doi: 10.1016/j.ijcard.2014.11.119.
61. Koizumi A, Sasano T, Kimura W, Miyamoto Y, Aiba T, Ishikawa T, et al. Genetic defects in a His-Purkinje system transcription factor, IRX3, cause lethal cardiac arrhythmias. *Eur Heart J*. 2016;37:1469–1475. doi: 10.1093/eurheartj/ehv449.
62. Visser M, van der Heijden JF, Doevendans PA, Loh P, Wilde AA, Hassink RJ. Idiopathic ventricular fibrillation: the struggle for definition, diagnosis, and follow-up. *Circ Arrhythm Electrophysiol*. 2016;9. doi: 10.1161/CIRCEP.115.003817.



63. Makita N, Behr E, Shimizu W, Horie M, Sunami A, Crotti L, et al. The E1784K mutation in SCN5A is associated with mixed clinical phenotype of type 3 long QT syndrome. *J Clin Invest*. 2008;118:2219–2229. doi: 10.1172/JCI34057.
64. Baruteau AE, Probst V, Abriel H. Inherited progressive cardiac conduction disorders. *Curr Opin Cardiol*. 2015;30:33–39. doi: 10.1097/HCO.000000000000134.
65. Smits JP, Veldkamp MW, Wilde AA. Mechanisms of inherited cardiac conduction disease. *Europace*. 2005;7:122–137. doi: 10.1016/j.eupc.2004.11.004.
66. Ingles J, Semsarian C. The value of cardiac genetic testing. *Trends Cardiovasc Med*. 2014;24:217–224. doi: 10.1016/j.tcm.2014.05.009.
67. Bamshad MJ, Ng SB, Bigham AW, Tabor HK, Emond MJ, Nickerson DA, et al. Exome sequencing as a tool for Mendelian disease gene discovery. *Nat Rev Genet*. 2011;12:745–755. doi: 10.1038/nrg3031.
68. Kapplinger JD, Giudicessi JR, Ye D, Tester DJ, Callis TE, Valdivia CR, et al. Enhanced classification of Brugada syndrome-associated and long-QT syndrome-associated genetic variants in the SCN5A-encoded Na(v)1.5 cardiac sodium channel. *Circ Cardiovasc Genet*. 2015;8:582–595. doi: 10.1161/CIRCGENETICS.114.000831.
69. Ackerman MJ. Genetic purgatory and the cardiac channelopathies: exposing the variants of uncertain/unknown significance issue. *Heart Rhythm*. 2015;12:2325–2331. doi: 10.1016/j.hrthm.2015.07.002.
70. Risgaard B, Jabbari R, Refsgaard L, Holst AG, Haunsø S, Sadjadih A, et al. High prevalence of genetic variants previously associated with Brugada syndrome in new exome data. *Clin Genet*. 2013;84:489–495. doi: 10.1111/cge.12126.
71. Refsgaard L, Holst AG, Sadjadih G, Haunsø S, Nielsen JB, Olesen MS. High prevalence of genetic variants previously associated with LQT syndrome in new exome data. *Eur J Hum Genet*. 2012;20:905–908. doi: 10.1038/ejhg.2012.23.
72. Ghouse J, Have CT, Weeke P, Bille Nielsen J, Ahlberg G, Balslev-Harder M, et al. Rare genetic variants previously associated with congenital forms of long QT syndrome have little or no effect on the QT interval. *Eur Heart J*. 2015;36:2523–2529. doi: 10.1093/eurheartj/ehv297.
73. Van Driest SL, Wells QS, Stallings S, Bush WS, Gordon A, Nickerson DA, et al. Association of arrhythmia-related genetic variants with phenotypes documented in electronic medical records. *JAMA*. 2016;315:47–57. doi: 10.1001/jama.2015.17701.
74. Maron BJ, Maron MS, Semsarian C. Genetics of hypertrophic cardiomyopathy after 20 years: clinical perspectives. *J Am Coll Cardiol*. 2012;60:705–715. doi: 10.1016/j.jacc.2012.02.068.
75. Ingles J, Semsarian C. Conveying a probabilistic genetic test result to families with an inherited heart disease. *Heart Rhythm*. 2014;11:1073–1078. doi: 10.1016/j.hrthm.2014.03.017.
76. Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, et al; ACMG Laboratory Quality Assurance Committee. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17:405–424. doi: 10.1038/gim.2015.30.
77. Rehm HL, Berg JS, Brooks LD, Bustamante CD, Evans JP, Landrum MJ, et al; ClinGen. ClinGen—the Clinical Genome Resource. *N Engl J Med*. 2015;372:2235–2242. doi: 10.1056/NEJMs1406261.
78. McNally EM, George AL Jr. New approaches to establish genetic causality. *Trends Cardiovasc Med*. 2015;25:646–652. doi: 10.1016/j.tcm.2015.02.013.
79. Chin D, Means AR. Calmodulin: a prototypical calcium sensor. *Trends Cell Biol*. 2000;10:322–328.
80. Shamgar L, Ma L, Schmitt N, Haitin Y, Peretz A, Wiener R, et al. Calmodulin is essential for cardiac IKS channel gating and assembly: impaired function in long-QT mutations. *Circ Res*. 2006;98:1055–1063. doi: 10.1161/01.RES.0000218979.40770.69.
81. Maier LS, Bers DM, Brown JH. Calmodulin and Ca<sup>2+</sup>/calmodulin kinases in the heart - physiology and pathophysiology. *Cardiovasc Res*. 2007;73:629–630. doi: 10.1016/j.cardiores.2007.01.005.
82. Alseikhan BA, DeMaria CD, Colecraft HM, Yue DT. Engineered calmodulins reveal the unexpected eminence of Ca<sup>2+</sup> channel inactivation in controlling heart excitation. *Proc Natl Acad Sci USA*. 2002;99:17185–17190. doi: 10.1073/pnas.262372999.
83. Fischer R, Koller M, Flura M, Mathews S, Strehler-Page MA, Krebs J, et al. Multiple divergent mRNAs code for a single human calmodulin. *J Biol Chem*. 1988;263:17055–17062.
84. Crotti L, Johnson CN, Graf E, De Ferrari GM, Cuneo BF, Ovidia M, et al. Calmodulin mutations associated with recurrent cardiac arrest in infants. *Circulation*. 2013;127:1009–1017. doi: 10.1161/CIRCULATIONAHA.112.001216.
85. Nyegaard M, Overgaard MT, Søndergaard MT, Vranas M, Behr ER, Hildebrandt LL, et al. Mutations in calmodulin cause ventricular tachycardia and sudden cardiac death. *Am J Hum Genet*. 2012;91:703–712. doi: 10.1016/j.ajhg.2012.08.015.
86. Boczek NJ, Gomez-Hurtado N, Ye D, Calvert ML, Tester DJ, Kryshtal DO, et al. Spectrum and prevalence of CALM1-, CALM2-, and CALM3-encoded calmodulin variants in long QT syndrome and functional characterization of a novel long QT syndrome-associated calmodulin missense variant, E141G. *Circ Cardiovasc Genet*. 2016;9:136–146. doi: 10.1161/CIRCGENETICS.115.001323.
87. Roux-Buisson N, Cacheux M, Fourest-Lieuvain A, Fauconnier J, Brocard J, Denjoy I, et al. Absence of triadin, a protein of the calcium release complex, is responsible for cardiac arrhythmia with sudden death in human. *Hum Mol Genet*. 2012;21:2759–2767. doi: 10.1093/hmg/dds104.
88. Chopra N, Yang T, Asghari P, Moore ED, Huke S, Akin B, et al. Ablation of triadin causes loss of cardiac Ca<sup>2+</sup> release units, impaired excitation-contraction coupling, and cardiac arrhythmias. *Proc Natl Acad Sci USA*. 2009;106:7636–7641.
89. Altmann HM, Tester DJ, Will ML, Middha S, Evans JM, Eckloff BW, et al. Homozygous/compound heterozygous triadin mutations associated with autosomal-recessive long-QT syndrome and pediatric sudden cardiac arrest: elucidation of the triadin knockout syndrome. *Circulation*. 2015;131:2051–2060. doi: 10.1161/CIRCULATIONAHA.115.015397.
90. Denegri M, Avelino-Cruz JE, Boncompagni S, De Simone SA, Auricchio A, Villani L, et al. Viral gene transfer rescues arrhythmogenic phenotype and ultrastructural abnormalities in adult calsequestrin-null mice with inherited arrhythmias. *Circ Res*. 2012;110:663–668. doi: 10.1161/CIRCRESAHA.111.263939.

KEY WORDS: inherited arrhythmia syndromes ■ pathogenicity ■ triadin, calmodulin

**New Insights Into the Genetic Basis of Inherited Arrhythmia Syndromes**  
Belinda Gray and Elijah R. Behr

*Circ Cardiovasc Genet.* 2016;9:569-577  
doi: 10.1161/CIRCGENETICS.116.001571

*Circulation: Cardiovascular Genetics* is published by the American Heart Association, 7272 Greenville Avenue,  
Dallas, TX 75231

Copyright © 2016 American Heart Association, Inc. All rights reserved.  
Print ISSN: 1942-325X. Online ISSN: 1942-3268

The online version of this article, along with updated information and services, is located on the  
World Wide Web at:

<http://circgenetics.ahajournals.org/content/9/6/569>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation: Cardiovascular Genetics* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Circulation: Cardiovascular Genetics* is online at:  
<http://circgenetics.ahajournals.org/subscriptions/>